

## DIRECTIONS

**RESRATTUS** is a complete WARFARIN rodenticide as received. Nothing need be added.

**RATS:** Effective use of RESRATTUS depends upon providing sufficient bait for successive feedings for ten or more days or until all signs of feeding have stopped. Place 4 ounces to 1 pound of bait in feeding stations located at intervals along rat runways or feeding areas. Inspect bait every day or two for the first week, and at regular intervals thereafter. Replenish baits as they are eaten, and replace if they become moldy or sour. When reinfestation is likely to occur, permanent bait stations should be established to guard against a build-up of the rat population.

**MICE:** Follow same directions for rats except that 1 or 2 tablespoons of RESRATTUS per station should be sufficient. Mouse control requires more feeding stations and more time than rat control. Place bait stations at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 10 days or until all signs of feeding have stopped.

NET CONTENTS: 1 POUND

# RESIDEX

## RESRATTUS

### KILLS RATS AND MICE WITH ONE FEEDING

ACTIVE INGREDIENTS	
Warfarin*	0.025%
INERT INGREDIENTS:	99.975%
Total	100%
*3-(4-acetylbenzyl)-4-hydroxy coumarin.	
U.S.D.A. Reg. No. 373-9	

**CAUTION:** Keep Out of Reach of Children  
(See Side Panel for Additional Caution Statements)

## PROFESSIONAL FORMULATION

MANUFACTURED BY  
**RESIDEX**  
CORPORATION  
225 TERMINAL AVENUE • CLARK, N. J.

4-3-70  
373-9

## PROPERTIES

RESRATTUS contains WARFARIN, a powerful blood anticoagulant which is readily accepted by rodents at effective dosage levels. Highly successful rodent control has resulted when RESRATTUS has been used according to directions. Rodent death results from internal hemorrhages if small quantities of RESRATTUS are taken over several consecutive days. Single isolated doses will not give the desired results.

## CAUTION

Follow directions carefully. Place RESRATTUS in bait boxes so it will not be exposed to humans, pets, poultry or livestock. In case baits are eaten accidentally, call physician at once. Note for Physician: When a human is known to have ingested RESRATTUS, blood transfusions along with oral and intravenous injections of Vitamin K are indicated as in the case of hemorrhages due to overdoses of Dicumarol.

NET CONTENTS: 1 POUND

# RESIDEX

## RESRATTUS

### KILLS RATS AND MICE WITH ONE FEEDING

ACTIVE INGREDIENTS	
Warfarin*	0.025%
INERT INGREDIENTS:	99.975%
Total	100%
*3-(4-acetylbenzyl)-4-hydroxy coumarin.	
U.S.D.A. Reg. No. 373-9	

**CAUTION:** Keep Out of Reach of Children  
(See Side Panel for Additional Caution Statements)

## PROFESSIONAL FORMULATION

MANUFACTURED BY  
**RESIDEX**  
CORPORATION  
225 TERMINAL AVENUE • CLARK, N. J.

## DIRECTIONS

RESRATTUS is a complete WARFARIN rodenticide as received. Nothing need be added.

**RATS:** Effective use of RESRATTUS depends upon providing sufficient bait for successive feedings for ten or more days or until all signs of feeding have stopped. Place 4 ounces to 1 pound of bait in feeding stations located at intervals along rat runways or feeding areas. Inspect bait every day or two for the first week, and at regular intervals thereafter. Replenish baits as they are eaten, and replace if they become moldy or sour. When reinfestation is likely to occur, permanent bait stations should be established to guard against a build-up of the rat population.

**MICE:** Follow same directions for rats except that 1 or 2 tablespoons of RESRATTUS per station should be sufficient. Mouse control requires more feeding stations and more time than rat control. Place bait stations at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days or until all signs of feeding have stopped.

NET CONTENTS: 1 POUND

# RESIDEX

## RESRATTUS KILLS RATS AND MICE with WARFARIN


ACTIVE INGREDIENTS	
Warfarin*	0.025%
INERT INGREDIENTS:	99.975%
Total	100%

\*3 (a acetonilbenzyl)-4-hydroxy coumarin.  
U.S.D.A. Reg. No. 373-9

**CAUTION:** Keep Out of Reach of Children  
(See Side Panel for Additional Caution Statements)

### PROFESSIONAL FORMULATION

MANUFACTURED BY  
**RESIDEX**  
CORPORATION  
225 TERMINAL AVENUE • CLARK, N. J.



## PROPERTIES

RESRATTUS contains WARFARIN, a powerful blood anticoagulant which is readily accepted by rodents at effective dosage levels. Highly successful rodent control has resulted when RESRATTUS has been used according to directions. Rodent death results from internal hemorrhages if small quantities of RESRATTUS are taken over several consecutive days. Single isolated doses will not give the desired results.

## CAUTION

Follow directions carefully. Place RESRATTUS in bait boxes so it will not be exposed to humans, pets, poultry or livestock. In case baits are eaten accidentally, call physician at once. Note for Physician: When a human is known to have ingested RESRATTUS, blood transfusions along with oral and intravenous injections of Vitamin K are indicated as in the case of hemorrhages due to overdoses of Dicumarol.

NET CONTENTS: 1 POUND

# RESIDEX

## RESRATTUS KILLS RATS AND MICE with WARFARIN

ACTIVE INGREDIENTS	
Warfarin*	0.025%
INERT INGREDIENTS:	99.975%
Total	100%

\*3 (a acetonilbenzyl)-4-hydroxy coumarin.  
U.S.D.A. Reg. No. 373-9

**CAUTION:** Keep Out of Reach of Children  
(See Side Panel for Additional Caution Statements)

### PROFESSIONAL FORMULATION

MANUFACTURED BY  
**RESIDEX**  
CORPORATION  
225 TERMINAL AVENUE • CLARK, N. J.