

ACCEPTED
 OCT 17 1983
 [Signature]

RESCUE™

POSTEMERGENT HERBICIDE

FOR SOYBEANS



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
WARNING
 Causes eye irritation. Do not get in eyes. Harmful if inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin or clothing.
ENVIRONMENTAL HAZARDS
 Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather favors drift from treated area

DIRECTIONS FOR USE
 It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage and disposal.
STORAGE: RESCUE is normally a solution but it may crystallize in the cold (32 F). Gentle warming (60 F) will redissolve crystals. Mild agitation such as rolling, shaking, or stirring before use is recommended.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
EQUIPMENT CLEANING: Thoroughly rinse spray tank, hose and nozzles with water after application. Use RESCUE in metal or fiberglass tanks. RESCUE can be removed from metal or fiberglass tanks by washing thoroughly with ammonia water and rinsing several times with water containing a good detergent.

COMPOSITION

Active Ingredient (percent by weight)	
Sodium salt of naptalam (N-1-Naphthylphthalamic acid)*	23.7%
Sodium salt of 4-(2,4-dichlorophenoxy) butyric acid**	6%
Inert Ingredients	75.7%
TOTAL	100.0

*Equivalent to 2 lbs. of the sodium salt of naptalam per gallon
 **Equivalent to 0.06 lbs 4-(2,4-DB) acid per gallon

KEEP OUT OF REACH OF CHILDREN
WARNING
STATEMENT OF PRACTICAL TREATMENT
 IF SWALLOWED: Drink two large glasses of water and induce vomiting. Contact a physician.
 IF ON SKIN: Wash affected area with plenty of soap and water.
 IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Contact a physician.
 IF INHALED: Remove to fresh air; if abnormal symptoms develop, contact a physician.
 SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Mfg by UNIROYAL CHEMICAL — Division of UNIROYAL, Inc
 Naugatuck, Connecticut 06770
 EPA REG. NO. 400-166
 EPA EST. NO.
 RESCUE™ is a trademark of Uniroyal, Inc
 Copyright 1983, Uniroyal, Inc

8 L 83
 Lot No.

NET CONTENTS

BEST

USE DIRECTIONS

GENERAL

RESCUE is a selective herbicide recommended for mid-to-late season postemergence weed control in soybeans. When applied according to label instructions, RESCUE will control or suppress the following broadleaf weeds, reducing their competition to the soybean crop: *cocklebur*, *giant ragweed*, *sunflower (wild & volunteer)*, *annual morningglory*, and *marshelder*.

Read and carefully observe all Application Notes prior to applying RESCUE.

APPLICATION TIMING

Determinate Soybeans - Apply when soybeans are at least 14 inches or taller.

Indeterminate Soybeans - Apply when soybeans are approximately 18 inches tall or after first bloom.

GROUND APPLICATION

Apply 2-3 quarts* of RESCUE in 10-25 gallons of water per acre (broadcast rate). A non-ionic spray adjuvant or crop oil concentrate at the rate of 2 quarts per 100 gallons of final spray solution should be used. The spray boom should be positioned 18-24" above the tops of the soybean plants or weeds and be equipped with hollow cone nozzles. Maintain a high pressure (40-50 p.s.i.) during application to insure better overall coverage. When making band applications, reduce rates and water volume according to width of band row spacing.

AERIAL APPLICATION

Apply 2-3 quarts* of RESCUE in a minimum of 5 gallons of water per acre. A non-ionic spray adjuvant or crop oil concentrate at the rate of 2 quarts per 100 gallons of final spray solution should be used. Nozzle size and orientation should provide maximum droplet breakup without causing excessive drift.

*For best results, use the lower rate of RESCUE (2 quarts) on weeds 10 to 12 inches or less in height and the higher rate (3 quarts) if weeds are taller than 12 inches. Adequate control or suppression can only be expected if application is made before weeds reach 3 feet in height.

APPLICATION NOTES

1. Always use a non-ionic spray adjuvant (80-20) or crop oil concentrate at the rate of 2 quarts per 100 gallons of final spray solution.
2. Avoid applications to drought, heat or disease stressed soybeans.
3. If rain occurs within 6 hours after application effectiveness may be reduced.
4. Drift-Applications should not be made when weather conditions favor drift. Do not apply by air within 50 feet of susceptible crops such as cotton, tomatoes or tobacco.
5. Soybean tolerance - Most soybeans are tolerant to RESCUE when used according to label instructions. Some leaf wrinkling or epinasty may occur under certain conditions, however, soybeans will usually outgrow this condition and continue to develop normally. Mitchell and Forrest varieties are more sensitive to RESCUE than other varieties.
6. DO NOT GRAZE OR FEED FORAGE FROM TREATED SOYBEANS TO LIVESTOCK.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.