

20 MULE TEAM[®]
MAINTAINA

For retarding growth of woody plants on
noncropland such as utility and highway rights
of way, parks and similar areas.

IMPORTANT

CARE SHOULD BE TAKEN TO CONFINE THE USE OR APPLICATION TO THE PARTS OF AN AREA INTENDED TO BE TREATED. KEEP PESTICIDE OFF TREATED AREA. DO NOT FEED TREATED VEGETATION TO HERD ANIMALS. DO NOT TREAT WOODY PLANTS USED FOR FEEDING ANIMALS.

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL

For use only by arborists, tree trimmers, etc.

As a retardant, often used as an effective dressing on cut surfaces in tree trimming practices. MAINTAINA controls a growth habit by controlling growth at an exposed trunk and inhibits sprouts from the cut surface area of the exposed area and also inhibits adventitious buds from developing near the treated cut. MAINTAINA has effectively retarded sprouting and/or eliminated branching in the following species:

American Elm	Tilia americana	Frst Oak	Quercus stellata
Bald Cypress	Taxodium distichum	Red Oak	Quercus rubra
Beech	Fagus spp.	Scot Cedar	Tamaria gallica
Birch	Betula spp.	Silver Maple	Acer saccharinum
Black Locust	Robinia pseudoacacia	Sycamore	Platanus spp.
Boxelder	Acer negundo	White Oak	Quercus nigra
Crabapple	Malus spp.	Wild Cherry	Prunus emarginata
Cypress	Cupressus spp.	Yellow Pine	Pinus ponderosa
Elm	Ulmus spp.	Quercus spp.	Quercus virginiana

DIRECTIONS

For wood control: Cut the tree. Apply with stiff brush or spray over cut surface including the exposed trunk and any other part of the bark and the underside of limb container after use.

NOTICE

The use of this product may result in killing shrubs and small trees. As protection seasonal weather, soil type, plant species, weight of cut, and temperature will affect results of these and other conditions may impair effectiveness. The use of this product may result in incomplete growth or browning of foliage. It is not guaranteed to be safe for use on State forests or woodlands. Agricultural Experiment Station weed control programs neither can nor do they guarantee that it will not result in any injury or damage to any plant material. The user of this product is advised to use it at his own risk. It is not recommended for use if there is any chance of it coming in contact with any animal. There is no express warranty or guarantee of any kind. The manufacturer disclaims all risk of usage and does not warrant that the product will be safe or effective. Neither manufacturer nor seller shall be liable for any personal injury or property damage resulting from the use of this product. The manufacturer and seller shall not be liable for any damages arising from the use of this product. No representations or warranties are made of the safety of this product.

© 1981 U.S.B.C.C.

* Trademark of United States Brush & Chemical Corporation

U.S. Patent 3,598,584

1981/118103

PAINT

FOR USE ONLY BY PROFESSIONAL TREE TRIMMING CREWS

A Tree Wound Dressing and Sprout Growth Retardant

For retarding growth of woody plants on noncropland such as utility
and highway rights of way, parks, and similar areas.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. Do not induce vomiting. May cause irritation to eyes, nose, throat and skin. Avoid contact with skin and clothing. Wash hands, face, and clothing after use.

Do not leave container where children or animals may take internally.

Avoid contamination of food and feed

DIRECTIONS FOR USE

Apply vegetation of FOLIATE LEAVES to the product in a manner incongruous with its function.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

ACCEPTED

OCT 15 1987

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

LEFFINGWELL

UNIROYAL
CHEMICAL

UNIROYAL CHEMICAL COMPANY INC

Middletown CT 06749

EPA REG NO 100-397

EPA EST NO

MAINTAIN is a Registered Trademark of UNIROYAL CHEMICAL COMPANY INC

NET CONTENTS ONE U.S. GALLON

8/21/87

Int No

USE DIRECTIONS

Stir well before using. Do not thin. Apply with a stiff paint brush. Uniformly cover cut surface including the exposed cambium and about one inch of the bark around the cut surface.

Store about 40°F. close container after use.

CHEMIGATION: Do not apply this product through any type of irrigation system

INSTRUCTIONS

Asphalt has long been used as an effective dressing on cut surfaces in tree trimming practices. MAINTAIN A combines a growth inhibitor, Chlorflurenol, with an asphalt emulsion and inhibits sprouts from the cut cambium layer of the exposed area and also inhibits adventitious buds from developing near the treated cut.


MAINTAIN A has effectively retarded sprouting and/or epicormic branching in the following species.

American elm	<i>Ulmus americana</i>
Bigtoothed aspen	<i>Populus grandidentata</i>
Black locust	<i>Robinia pseudacacia</i>
Black walnut	<i>Juglans nigra</i>
Box elder	<i>Acer negundo</i>
Decorative olive	<i>Oleaaceae spp</i>
Golden willow	<i>Salix lasiandra</i>
Hickory	<i>Carya spp</i>
Live oak	<i>Quercus virginiana</i>
Post oak	<i>Quercus stellata</i>
Red oak	<i>Quercus rubra</i>
Salt cedar	<i>Tamarix gallica</i>
Silver maple	<i>Acer saccharinum</i>
Sycamore	<i>Platanus spp.</i>
Water oak	<i>Quercus nigra</i>
Wild cherry	<i>Prunus emarginata</i>
Yellow poplar	<i>Liriodendron tulipifera</i>

IMPORTANT

Care should be taken to confine the use or application to the particular area intended to be treated. Keep livestock off treated area. Do not feed treated vegetation to livestock. Do not reuse container. Destroy when empty. Apply this product only as specified on this label.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



MAINTAIN™ A PAINT

FOR USE ONLY BY PROFESSIONAL TREE TRIMMING CREWS

A Tree Wound Dressing and Sprout Growth Retardant

For retarding growth of woody plants on noncropland such as utility
and highway rights of way, parks, and similar areas.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. Do not induce vomiting. May cause irritation to eyes, nose, throat and skin. Avoid contact with skin and clothing. Wash hands, face, and clothing after use.

Do not leave container where children or animals may take internally.

Avoid contamination of food and feed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

COMPOSITION

Active Ingredients by weight	
Methyl 2-chloro-9-hydroxyfluorene-9-carboxylate	5.13
Methyl 9-hydroxyfluorene-9-carboxylate	0.04
Methyl 2,7-dihydro-9-hydroxy-fluorene-9-carboxylate	0.03
INERT INGREDIENTS	59.75
Total	100.00

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

LEFFINGWELL™

**UNIROYAL
CHEMICAL**

1000 LEFFINGWELL STREET, MIDDLEBURY, CONNECTICUT 06762

UNIROYAL CHEMICAL COMPANY INC

Middlebury, CT 06749

EPA REG. NO. 400-397

EPA EST. NO.

MAINTAIN is a Registered Trademark of UNIROYAL CHEMICAL COMPANY INC

NET CONTENTS ONE U.S. GALLON

B21 R1

Ed. No.