

[407-310]

**IMPERIAL
PYRENONE ALL-PURPOSE SPRAY**

DIRECTIONS FOR USE

Hold container upright, nozzle away from you. Press valve down and spray as directed.

**FOR THE CONTROL OF
HOUSEHOLD INSECTS**

Do not remain in treated areas, and ventilate the area after treatment is completed. To control house flies, mosquitoes, moths, and gnats, shut doors and windows and spray room at the rate of 5 to 6 seconds per 1,000 cubic feet. Keep room closed for at least 15 minutes after spraying.

To kill roaches and silverfish: Spray directly into cracks, crevices, and other secluded places. Direct spray at insects when they appear.

To kill ants: Spray directly on insects and any trails and other places where ants congregate.

**FOR THE CONTROL OF
ORNAMENTAL GARDEN INSECTS**

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Do not wet foliage or blossoms.

For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray, as well as aphids, exposed thrips, spider mites, lace bugs, clover mites and army worms.


May be used on plants, as a multi-purpose spray in small ornamental gardens, and for spot treatment of incipient insect infestations to control spreading in large ornamental garden areas.

May also be used on other plants, such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering jew and zinnias.

FOR USE AS AN AREA SPRAY

This new formula double-purpose insect killer is designed for use, indoors and outdoors, to kill flies, mosquitoes, moths and gnats. A general misting of the area, when the air is quiet, should be sufficient to reduce the number of these pests during meals. If flies or mosquitoes tend to congregate in a given area, it should be sprayed directly upon them. For control, spray as often as necessary during outdoor barbecuing. May be sprayed on the grill and tables, but do not spray directly on food.

To Treat Cattle and Horses: To protect cattle and horses from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray the eyes. Repeat treatment as necessary when flies are troublesome.



All-Purpose Spray

**FOR RAPID CONTROL OF FLIES, MOSQUITOES
AND GNATS IN HOMES, ANIMAL QUARTERS,
MILK ROOMS AND ON LIVESTOCK**

**Kills stable flies, horse flies, face flies, deer flies,
house flies, horn flies, mosquitoes and gnats.**

**Designed for Use in Edible Product Areas of
Official Establishments Operating Under the
Meat, Poultry, Shell Egg Grading and Egg
Products Inspection Programs, and in Other
Food Processing Plants.**

ACTIVE INGREDIENTS:	
Pyrethrins	0.5%
Piperonyl Butoxide, Technical*	5.0%
Petroleum Distillate	2.0%
INERT INGREDIENTS:	92.5%
	100.0%

*Equivalent to 4.0% (butylcarbityl) (8-propylpiperonyl) ether and to 1.0% of related compounds.

PYRENONE is a registered trademark of FMC Corp.

**CAUTION
KEEP OUT OF REACH
OF CHILDREN**

(See back panel for additional precautions)

NET WT. 16 OZ. AVD.

**Manufactured for
IMPERIAL INC.
SHENANDOAH, IOWA 51601**

**IMPERIAL
PYRENONE ALL-PURPOSE SPRAY**

DIRECTIONS FOR USE

In Barns, Dairy Barns, Stables and Milk Rooms: For rapid control of flies, mosquitoes, moths and gnats close windows and doors and spray at the rate of five to six seconds per 1,000 cubic feet of space. For best results keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

Food Plants: Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

Close windows and doors and shut off ventilating system. Apply at the rate of five to six seconds per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, mosquitoes, flying moths, gnats, wasps and hornets.

Direct application to exposed insects will control the following stored product pests: grainary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, mealworms, and grain mites and these common pests: silverfish, crickets, roaches, spiders, scorpions, cheese skippers and clover mites.

CAUTION: Harmful if swallowed. Avoid breathing mist.

Remove pets, birds and cover fish aquariums before spraying.

WARNING: Contents Under Pressure. Do not puncture container nor store at over 120 F. Do not incinerate. Do not use in areas where fish or reptiles may be exposed to the spray. Do not apply over uncovered milk or food products. Any exposed utensil or food handling surfaces should be thoroughly cleaned before use.

Do not reuse container. Dispose of by burying with waste.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 107-310

M-2363B-L