

[419-90]

DIRECTIONS:

HORN FLIES ON BEEF AND DAIRY CATTLE: Apply 2 ounces (4 table-spoons) of 4% Malathion Dust per animal. Apply dust on back and neck of animal and repeat application at 10 to 14 day intervals. Do not apply to dairy cattle later than five hours before milking. Ideal time for application is immediately after milking. Do not use spray or dip dairy animals in Malathion. Do not contaminate water troughs. Do not use on calves less than one month old.

DIRECT APPLICATION TO CHICKENS, DUCKS, GEESE, AND TURKEYS: For the control of Northern Fowl Mite, Poultry Lice and as a supplement to premise treatment for Chicken Red Mite, dust individual birds. Apply dust by means of a shaker can or with a rotary hand duster. Repeat in 4 to 8 weeks or when necessary.

POULTRY HOUSES: Apply 1 lb. of dust per 50 to 60 square feet of litter and floor space and to nests, roosts and adjacent areas. Replace treated nests with clean, new nesting material. Dust may be applied by sprinkling from a can, or with a rotary hand duster or puff duster. Repeat as necessary.

FOR LICE ON BEEF CATTLE AND NON-LACTATING DAIRY CATTLE: Apply thoroughly to animals. Repeat application in 10 days and thereafter as necessary. Do not use on non-lactating dairy animals within two weeks of freshening. Do not use on calves less than one month old.

FOR LICE ON SWINE: Apply thoroughly to animals. Pens should also be thoroughly dusted. Repeat in 10 days and thereafter as needed.

CAUTION: Avoid breathing dust and prolonged and repeated contact with skin. Wash hands after use. Avoid contamination of feed, feeding troughs, and water fountains. Use on dairy animals only in accordance with directions. Keep out of reach of children.

Concol

**4%
MALATHION
DUST**

**FOR ON DAIRY CA
CATTLE,
POULTRY
and
SWINE**

2419-90

U.S.D.A. REG. NO. 419-90

DIRECTIONS:

HORN FLIES ON BEEF AND DAIRY CATTLE: Apply 2 ounces (4 table-
spoons) of 4% Malathion Dust per animal. Apply dust on back and
neck of animal and repeat application at 10 to 14 day intervals. Do
not apply to dairy cattle later than five hours before milking. Ideal
time for application is immediately after milking. Do not use spray
or dip dairy animals in Malathion. Do not contaminate water troughs.
Do not use on calves less than one month old.

DIRECT APPLICATION TO CHICKENS, DUCKS, GEESE, AND TURKEYS:
For the control of Northern Fowl Mite, Poultry Lice and as a supple-
ment to premise treatment for Chicken Red Mite, dust individual birds.
Apply dust by means of a shaker can or with a rotary hand duster.
Repeat in 4 to 8 weeks or when necessary.

POULTRY HOUSES: Apply 1 lb. of dust per 50 to 60 square feet of
litter and floor space and to nests, roosts and adjacent areas. Replace
treated nests with clean, new nesting material. Dust may be applied
by sprinkling from a can, or with a rotary hand duster or puff duster.
Repeat as necessary.

FOR LICE ON BEEF CATTLE AND NON-LACTATING DAIRY CATTLE:
Apply thoroughly to animal's. Repeat application in 10 days and
thereafter as necessary. Do not use on non-lactating dairy animals
within two weeks of freshening. Do not use on calves less than one
month old.

FOR LICE ON SWINE: Apply thoroughly to animals. Pens should also
be thoroughly dusted. Repeat in 10 days and thereafter as needed.

CAUTION: Avoid breathing dust and prolonged and repeated contact
with skin. Wash hands after use. Avoid contamination of feed, feed-
ing troughs, and water fountains. Use on dairy animals only in
accordance with directions. Keep out of reach of children.

CAUTION: Do not reuse empty container. Destroy it by burying with
waste or burning. Stay away from smoke or fumes.

This product is toxic to fish. Keep out of lakes, streams or ponds.
Do not contaminate water by cleaning of equipment, or disposal of
wastes. Apply this product only as specified on this label.

Manufactured by
CENOL COMPANY, INC. • CHICAGO, ILL. 60651

ACCEPTED

July 2, 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTE
ED UNDER NO. 419-90 SUB
TO ATTACHED COMMENTS.

**4%
MALATHION
DUST**

**FOR USE
ON DAIRY CATTLE**
**FOR
CATTLE,
POULTRY
and
SWINE**

CAUTION —Keep out of reach of children.
See other warnings on back panel.

ACTIVE INGREDIENTS:

Malathion—O, O dimethyl dithi-
ophosphate of diethyl mercapto-
succinate 4.0%

INERT INGREDIENTS 96.0%

NET CONTENTS 1 POUND