

DIRECTIONS FOR USE

Application should be made only by experienced personnel using approved safety equipment. Apply when air is calm, using ground equipment or airplane. Make coverage coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control. DO NOT EXCEED THE LIMITS GIVEN FOR SPECIFIC CROPS.

RECOMMENDATIONS

BEANS (Bush and Pole): Mexican Bean Beetle, Thrips--use 15 to 25 lb. per acre. Serpentine Leaf Miner--use 20 to 25 lb. per acre. Stinkbugs--use 30 lb. per acre. Make no application to cattle or meat animals.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE: Aphids, Serpentine Leaf Miner--use 20 to 25 lb. per acre. Cabbage Webworm--use 15 to 25 lb. per acre. Do not apply on Broccoli and Brussels Sprouts after edible parts start to form. May be applied on Cabbage up to 7 days before harvest if outer leaves are to be stripped at harvest. If outer leaves are not stripped, do not apply after heads start to form.

EGGPLANT: Aphids, Red Spider Mites--use 30 lb. per acre. Thrips--use 15 to 20 lb. per acre. Serpentine Leaf Miner--use 20 lb. per acre. Do not apply within 15 days of harvest. Do not apply after edible parts start to form.

LETTUCE: Aphids--use 30 lb. per acre. Armyworms, Cutworms (climbing)--use 10 to 20 lb. per acre. Do not apply to leaf lettuce after seedling stage or to head lettuce after heads start to form. If outer leaves of head lettuce are stripped at harvest, application may be made up to 21 days of harvest.

TOMATOES: Aphids, Serpentine Leaf Miner--use 20 lb. per acre. Leaf-footed Plant Bug, Thrips, Tomato Hornworm--use 15 to 20 lb. per acre. Stinkbugs--use 30 lb. per acre. Do not apply within 10 days of harvest.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured by

WOODBURY CHEMICAL COMPANY

St. Joseph, Mo. Denver, Colo. Orlando, Fla. Lubbock, Tex.

F.A. Reg. No. 441-