

ACCF T D
 5-12-67
 421-17
 UNDER THE FEDERAL REGISTER AND FOR EXPORT TO UNDESIGNED TO ATTACHED

DIRECTIONS FOR USE

FOR THE SICK ROOM: All utensils, such as bed-pans, urinals, etc., wash with a solution of three ounces of disinfectant to 1/2 gallon of water.

For **BOILING CLOTHING** and **LINEN** used in sick room: Use in the proportion of three ounces of disinfectant to one gallon of water. Add the water to the disinfectant and mix well before using.

WATER CLOSETS and **URINALS.** Use 3 ounces of disinfectant to a gallon of water. Apply freely to all surfaces to be disinfected and scrub the exposed areas.

FOR GENERAL DISINFECTING

Remove all filth and dirt from the premises and use a solution of one part disinfectant to 100 parts of water. This mixture acts as a disinfectant against **S. typhosa**

In the typhoid sick room, in the disinfecting of beds, clothes, utensils, typhoid stools, rooms, and the hands of those attending typhoid patients, disinfectant should be used in the proportion of 4 ounces to each gallon of water.

Use one-half cup of the disinfectant to two gallons of water for mopping, scrubbing, sprinkling or cleaning purposes.

GARBAGE: To deodorize, sprinkle with solution, one-half cup to a gallon of water. Wash the empty can, with some of the solution and leave the wash in can.

NET CONTENTS:

**ODORLESS
 DISINFECTANT**

Phenol Coefficient at least 5 by ^{A.O.A.C.} Method

ACTIVE INGREDIENTS 10%

1. VEGETABLE OIL SOAP	3. ORTHO-PHENYL PHENOL - 1 word
2. CHLORO-2-PHENYL PHENOL	4. ISOPROPYL ALCOHOL

INERT INGREDIENT Water 90%

Preferably add water to the disinfectant

~~FOR ANIMAL QUARTERS~~ tion of disinfectant in the proportion of one cupful to a pail of water. Scrub all floorings with same solution. This will help prevent disease.

14 pt

FOR EXTERNAL USE ONLY

CAUTION: KEEP OUT OF REACH OF CHILDREN
 (See side panel for further caution)

Manufactured by
 James Varley & Sons, Inc. St. Louis, Missouri

10 pt

DIRECTIONS FOR USE

~~Red Mites: Paint or spray the perches, nest boxes, drop boards, floor and inside walls, poultry houses and coops with full strength disinfectant, repeating as often as necessary. Applied in this manner, it is harmless to the birds but should not be used directly on the fowl.~~

This disinfectant can be used anywhere where clean and sanitary quarters are desired. A partial list of places using this type of disinfectant follows: Hotels, schools, institutions, office buildings, factories, jails, public buildings, railroad terminals, theatres, stores, sanitariums and many others.

~~This disinfectant has certain advantages over coal tar and cresol disinfectants in that it is relatively non-irritating, non-corrosive and does not injure cloth, metals, etc., does not injure body tissues, is comparatively cheap in price, is not affected appreciably by organic matter and has a pleasant odor.~~

Every batch is tested for germicidal strength in a modern bacteriological laboratory - but before these tests are made the raw materials are checked in our chemical laboratories - and the finished product is then double checked.

CAUTION: May cause skin irritation.
 Avoid contact with skin and eyes. In case of contact with skin, wash with water.

USDA Reg. No. 421-17