

PROPOXUR TRANSPARENT EMULSION** PRESSURIZED SPRAY 0.5%

Offering Good Flushing Activity and Residual Kill of Cockroaches for 2 to 3 Weeks

FOR RESIDUAL CONTROL OF ANTS, COCKROACHES (WATERBUGS), CLOVER MITES, CRICKETS, EARWIGS, FLEAS, MOSQUITOES (OUTDOORS), HOUSE FLIES (OUTDOORS), HORNETS, MILLIPEDES, CENTIPEDES, SAW-TOOTHED GRAIN BEETLES (EXPOSED STAGE), SCORPIONS, SILVERFISH (FIREBRATS), SOWBUGS (PILLBUGS), SPIDERS, BROWN DOG TICKS, AND WASPS

ACCEPTED
 JUL 20 1983
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 432-699

ACTIVE INGREDIENTS:

o-isopropoxyphenyl methylcarbamate

0.5%w/w

INERT INGREDIENTS:

99.5%

100.0%w/w

** U.S. Patent No. 3,683,078

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que se explique a usted en su idioma. (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

In food handling establishments, can be used in non-food areas as a contact or residual spray. In food areas, application is limited to crack and crevice treatment.

SHAKE WELL BEFORE USING

EPA Est.: See Lot No. For Correlation

EPA Reg. No. 432-699

G 42545-MO-1

P-5/10/88

H 5905-AR-1

P 655-GA-1

NET WEIGHT:

ROUSSEL BIO
 CORPORATION
 PO Box 1077, 400 Sylvan Avenue
 Englewood Cliffs, New Jersey 07632

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if inhaled. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing.

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not apply to animals or humans. Do not use on food crops. Do not use in animal barns or poultry houses for fly control. In food handling areas, application is limited to crack and crevice and/or spot treatment. Do not apply directly to food and feedstuffs. In commercial food handling facilities, cover or remove any food or food processing equipment during application. Do not apply while food processing is underway. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water.

If on skin: Wash with soap and water.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal only if symptoms of cholinesterase inhibition are present.

PHYSICAL & CHEMICAL HAZARDS

Contains under pressure. Do not puncture or incinerate container. Do not use near heat or open flame. Exposure to temperatures above 130°F (54.5°C) may cause bursting.

Do not apply this water-based product in conduits, motor housings, junctions and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be stained or harmed by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep in a locked storage area.
CONTAINER DISPOSAL: Replace cap and discard container in trash.
 Do not incinerate or puncture.

DIRECTIONS FOR USE (Continued on reverse side)

M/

Shake well before each use. Remove protective cap. Hold can upright and aim spray opening away from person. Push button to spray.

INDOORS: Pest control in and around homes, apartment buildings, stores and warehouses: For residual control of ants, cockroaches (waterbugs), millipedes, centipedes, silverfish (firebrats), spiders, fleas, brown dog ticks and to aid in control of saw-toothed grain beetles (exposed stage). Apply spray until area is wet but not to the point of run-off. Use as residual spray, hitting as many insects as possible, to baseboards, window frames, the undersides of shelves and drawers, under sinks and stoves and to other places where insects may hide. Also spray cracks, crevices and surfaces where insects may crawl when they come out of hiding. For ants, spray trails and places where ants enter premises. Repeat as necessary.

Do not spray foodstuffs or house plants. Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated surfaces until surfaces are completely dry.

Pest control in non-food areas only of meat packaging and food processing plants, food storage areas, restaurants, and other food handling establishments (places other than private residences in which exposed food is held, processed, prepared or served). Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). For residual control of ants, cockroaches, millipedes, silverfish, spiders, waterbugs, and to aid in control of saw-toothed grain beetles (exposed stage).

Apply to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

FOOD AREAS: Limited to crack and crevice treatment only. Apply a small amount of material as a pin-thin stream directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, or hollow equipment legs where the insects noted above hide. Do not use this product in conduits, motor housings and electrical switch boxes.

Care should be taken to avoid depositing the product onto exposed food processing surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

Applications of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

Spraying unloading docks and outside areas where insects congregate will aid in preventing reinfestations. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

Pest control in animal research facilities: The use of this product on and around research animal quarters for pest control will not induce changes in liver microsomal enzyme activity in these animals.

PESTS OUT-OF-DOORS
DO NOT TREAT OR ALLOW THE SPRAY TO CONTACT PLANTS (LAWNS, FLOWERING PLANTS, SHRUBS, TREES, EVERGREENS, ETC.) BECAUSE OF POSSIBLE INJURY

For residual control of insects around industrial buildings, homes, garbage areas, dumps.

RESIDUAL SPRAY for ants, cockroaches (waterbugs), earwigs, house flies, hornets, millipedes, centipedes, mosquitoes, spiders, crickets, clover mites, fleas, scorpions, sowbugs (pillbugs), brown dog ticks, and wasps. Apply spray until area is wet but not to the point of run-off. Use as a residual spray, hitting as many insects as possible, to surface of buildings, porches, screens, window frames, patios and garages where these insects congregate. For control of wasps and hornets, apply after dark when all of the insects have returned to the nest. Repeat as necessary.

Delete
JD

WARRANTY STATEMENT

Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with the directions for use or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

70