

A knapsack sprayer is suitable for small areas. For larger areas tractor attachments with one or more hoses are faster and more economical.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

**CAUTION
HARMFUL IF SWALLOWED**

Avoid contact with skin, eyes, or clothing. Do not contaminate water to be used for irrigation or domestic purposes. Avoid spray to susceptible plants as this may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use same spray equipment for other purposes). Do not store near fertilizers, seeds, insecticides, fungicides. Vapors from this product may cause injury to susceptible plants growing nearby. Do not use around the home, recreation area or similar areas.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy by burying in a safe place. Do not burn.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured For
TECHNE CORPORATION
St. Joseph, Mo. 64502; Denver, Colo. 80216

EPA Reg. No. 449-218

Techne

ACCE

December

UNDER THE FEDERAL
FUNGICIDE AND RO
FOR ECONOMIC RO
ED UNDER NO. 4

No. 6 LOW VOLATILE 2,4,5-T

CONTAINS 6 POUNDS 2,4,5-T ACID PER GALLON

ACTIVE INGREDIENTS:

*2,4,5-Trichlorophenoxyacetic Acid, 90.9%
Propylene Glycol Butyl Ether Ester

INERT INGREDIENTS

TOTAL

9.1%

100.0%

*Equivalent to 2,4,5-Trichlorophenoxyacetic acid 59.2%

CONTROLS most species of woody plants except Prickly Ash, Basswood, Black Birch and Scrub Oak. Some susceptible species may require 2 or more applications.

GENERAL DIRECTIONS

May be applied by airplane, ground rigs, or hand dispensers. This product may be used for Brambles, Hawthorn, Osage Orange, Blackjack or Post Oak, Persimmon, or Wild Rose in fence rows, right-of-ways, roads, vacant lots and other similar non-crop areas.

FOLIAGE SPRAY: Apply spray to both foliage and stems of all plants from time foliage reaches full maturity until plants begin to go dormant. In high volume sprayers mix 3 1/3 pints per 100 gallons of water per acre. In low volume sprayers, mix 3 1/3 pints in 5 to 10 gallons of water depending on type of equipment. For aerial application use 3 1/3 pints in 6 to 8 gallons of water. Aerial applicators may find conditions vary according to the density of foliage and the woody species present. It is suggested that they refer to State Agricultural Experiment Stations for best local recommendations.

BASAL SPRAY OR STUMP TREATMENT

Use 2 1/3 gallons in 100 gallon of oil (for smaller areas, use 10 oz. in 3 gallons of oil.) Apply this spray mixture to the lower 12 inches of the stem or trunk around the entire perimeter and to all exposed roots. It is important that spray be applied to the point of run-off.

The same procedure applies to stump application after cutting. The spray should cover the entire stump to the ground and be applied to the point of run-off.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL
CAUTIONS**