

GENERAL INFORMATION

It is best to prevent plant diseases by following a regular spray program. Start applications when vegetables are first set in field or garden, and for roses, when first leaves are fully formed in Spring. Make full coverage applications every 7 to 10 days throughout growing season. Shorten spray intervals when disease appears, or when weather conditions are conducive to spore activity (warm, wet and humid weather). When these conditions exist, make applications every 3 to 4 days.

For each gallon water to be used, add 2 level tablespoons of Chemform Maneb Fungicide. First make a paste by adding Chemform Maneb to small amount water, and then add paste to remaining water. 2 tablespoons to each gallon water will give adequate protection and control. For additional information for usage of Chemform Maneb Fungicide on other plants in your area, consult your county, state or regional Agricultural Agent or Extension Service.

Use any good type pressure, tank or garden hose sprayer. Make thorough coverage applications. Spray entire plant paying particular attention to under-side of leaves. For additional protection from diseases, spray immediate soil surface around plant, at time plant is sprayed, in order to kill disease spores that may come from soil. Good sanitation practices of removing diseased leaves, destroying dead vines, and a regular feeding program will also aid in controlling plant diseases.

NET. WT. 6 OZ.

MAKES 23 GALLONS
SPRAY FUNGICIDE



*Town 'n
Country
Chemicals*

"The Best for Gardens Best"

MANEB FUNGICIDE SPRAY POWDER

CAUTION: KEEP OUT OF REACH OF CHILDREN.

**CONTROLS ROSE BLACK SPOT
DISEASE AND CERTAIN DISEASES
OF TOMATOES AND POTATOES.**

U.S.D.A. Reg. No. 456-87

Active Ingredients:

MANEB	
(Manganese ethylenebisdithiocarbamate) . . .	40.0%
(Manganese expressed as metallic—16.5%)	
Inert Ingredients	60.0%
Total	100.0%

DIRECTIONS:

BLACK SPOT DISEASE of Roses: Start spray applications after new Spring leaves are well formed, and spray every 7 to 10 days throughout growing flowering season. During high humid wet weather conditions, conducive to disease spore activity, decrease days between sprays. Make thorough coverage sprays: both sides of leaves and stems. For additional protection and prevention of disease spores that may rise from surrounding soil—spray soil surface around plant at time plant application is made.

TOMATO: Early and Late Blights, Leaf Spot (Septoria), Gray Leaf Spot and Anthracnose: Spray plants every 7 to 10 days throughout growing season. Start spray schedule when plants are set in field or garden. Spray soil at time plant is sprayed for additional protection from disease. Do not use on seedling plants in cold-frames or greenhouses.

POTATO: Early and Late Blight: Start sprays when plants are 3 to 4 inches high and continue applications every 7 to 10 days until tops start yellowing at end of growing season.

NOTE: CHEMFORM'S MANEB FUNGICIDE SPRAY POWDER is compatible and may be used with many commonly used insecticides for combination-fungicide-insecticide control. For a better fungicide-insecticide combination, use CHEMFORM'S MANEB with CHEMFORM'S LIQUID BARICIDE Multi-Purpose insecticide Spray Concentrate. For MANEB-BARICIDE combination spray use 1 tablespoon of Liquid Baricide and 2 tablespoons Chemform's Maneb to each gallon of water to be used.

Caution: May cause irritation of eyes, throat, nose and skin. May be harmful if swallowed. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. In case of contact, flush with quantity of water; for eyes, get medical attention.

The use, storage and handling of this product being beyond the control of seller and manufacturer, neither makes any warranty, express or implied. Buyer assumes all risk of use, storage and handling, whether in accordance with directions or not.

**MANUFACTURED BY
CHEMICAL FORMULATORS, INC.
NITRO, W. VA.**