

50 LB.



APR 24, 1969

NET

# TOXAPHENE-SULFUR 20-40 DUST

## AN AGRICULTURAL INSECTICIDE - FUNGICIDE

### ACTIVE INGREDIENTS:

*Toxaphene .....	20.0%
Sulfur .....	40.0%
<b>INERT INGREDIENTS</b> .....	<b>40.0%</b>
*Technical Chlorinated Camphene (Chlorine Content 67% to 69%) .....	100.0%

### DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and infection and repeat at 5- to 10-day intervals, depending upon the severity of the disease and insect conditions, but observe use limits given for specific crops.

For the control of diseases persisting after the prescribed time limits have expired, an effective fungicide should be used alone, in a manner which will not leave residues in excess of established tolerances.

### CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED.  
 AVOID BREATHING DUST. AVOID CONTACT WITH EYES, SKIN, AND CLOTHING.  
 WASH THOROUGHLY AFTER HANDLING.  
 DO NOT CONTAMINATE FOOD AND FEED.

### OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.**

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. *Birds feeding on treated areas may be killed.*

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

Do not use Toxaphene or sulfur on cucurbits (cantaloupe, cucumber, squash, melon), or young plants, as injury to plants may result.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.

### RECOMMENDATIONS

**BEANS:** Armyworms, Bean Leaf Roller and Leafhopper, Climbing Cutworms (apply only in seedling stage), Corn Earworm, Cowpea Curculio, Lygus, Powdery Mildew, Rust, Salt-marsh Caterpillars, Thrips -- Use 15 lb. per acre. **Do not apply to green or snap beans within 7 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.**

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Cabbage Looper, Corn Earworm -- Use 10 to 20 lb. per acre. Imported Cabbageworm, Cutworms (climbing and surface-feeding), Fall Armyworm, Serpentine Leaf Miner, Thrips -- Use 10 lb. per acre. Flea Beetles -- Use 15 lb. per acre. ~~Do not apply to heads beginning to form.~~

**COTTON:** Overwintered Boll Weevils, Flea Beetles, Fleahoppers, Thrips (early-season control) -- Apply 10 lb. per acre. Make 2 to 4 applications at 7 to 10-day intervals. Cotton Leafworm, Fall Armyworm, Lygus and other Mirids -- Use 10 to 15 lb. per acre. Boll Weevil, Bollworm -- Use 15 to 20 lb. per acre. Apply at 4 to 5-day intervals until control is obtained. **Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.**

**PEPPERS:** Armyworms, Blister Beetle, Colorado Potato Beetle, Cutworm (climbing and surface-feeding), Flea Beetle, Hornworm, Serpentine Leaf Miner -- Use 10 to 15 lb. per acre. Fruitworm, Pepper Weevil -- Use 15 lb. per acre. **Do not apply within 5 days of harvest.**

**TOMATOES:** Armyworms, Blister Beetle, Cabbage Looper, Colorado Potato Beetle, Cutworms (climbing and surface-feeding), Flea Beetle, Pinworm, Russet Mite, Serpentine Leaf Miner, Tomato Fruitworm, Tomato Hornworm -- Use 10 to 25 lb. per acre. **Do not apply within 3 days of harvest.**

**NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.**

BURN OR BURY EMPTY CONTAINER — STAY OUT OF SMOKE

USDA Reg. No. 476-176  
 Made in U.S.A.  
 Stauffer Chemical Company, New York  
 T TA 680419 SE  
 690419

# STAUFFER CHEMICAL CO. NEW YORK