

HEPTACHLOR 10 GRANULAR

20-40 Mesh — U.S. Sieve Series Designation

FOR USE AS A SOIL INSECTICIDE

50 LB. NET

ACTIVE INGREDIENTS:

Heptachlor* 10%
Related Compounds 4%

INERT INGREDIENTS

*Heptachlorotetrahydro-4,7-methanoindene 86%
100%

WARNING

Keep Out of Reach of Children

- May be fatal if swallowed
- Do not breathe vapor or dust
- Do not get on skin
- In case of contact, wash immediately with soap and water
- Do not contaminate feed and food

USE PRECAUTIONS

In order that pesticidal residues on food and forage crops will not exceed maximums established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas accessible to unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

To protect bees, do not apply when crops are in bloom. Remove or cover bees, if applicable, during bloom.

Highly toxic to fish and wild life. Do not contaminate any body of water, water courses, or other non-crop areas.

Do not use this product on feed or forage to be fed to dairy animals or to animals being prepared for slaughter.

Do not use this product on feed or forage to be fed to dairy animals or to animals being prepared for slaughter.

DIRECTIONS FOR SOIL TREATMENT

BROADCAST: Make preplant or adjuvant application with a granular spreader or fertilizer spreader or by airplane application to cover this soil surface uniformly. Follow up with water to mix the insecticide into the top 3 to 6 inches of soil.

ROW OR BAND TREATMENT: Apply the recommended amount of Heptachlor 10 granular to the planting or later attachment to the row or planting zone.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

RECOMMENDATIONS

CROP	INSECT	
BERRIES (Blackberries, Raspberries, Elderberries, Huckleberries, Rubus spp., and other raspberries)	Citrus root weevil	40
CITRUS	Citrus root weevil	30
CORN	Wireworms, seed corn maggot, green June beetle grubs, white grubs, cutworms	10-20
	Wireworms	15
	Wireworms, seed corn maggot, green June beetle grubs, white grubs	20-30
PEPPERS, TOMATOES	Cutworm, seed corn maggot, wireworms	20
	Wireworms, white grubs	30
SMALL GRAINS (Barley, Oats, Rye and Wheat)	Wireworms, white grubs	20-30

FOR USE BY COMMERCIAL APPLICATORS ONLY

INSECT CONTROL ON ORNAMENTAL TURF, LAWN, NURSERY AND GREENHOUSE SOILS

ANTS	EUROPEAN CHAFER GRUBS	PILLBUGS
ARMYWORMS	GREEN JUNE BEETLE GRUBS	SNAILS
ASIATIC GARDEN BEETLE GRUBS	JAPANESE BEETLE GRUBS	SCORPIONS
CUTWORMS	LAWN CHINCH BUGS	STRIPED GRASSWORMS
EARTHWORMS	MILK CRICKETS	WHITE GRUBS
	NORTHERN MASKED CHAFER GRUBS	

For use in turf soils, as golf courses, athletic fields, lawns, ornamental gardens, cemeteries, and in seed and nursery and greenhouse soils where ornamental plants are grown, use 30 lbs. of Heptachlor 10 granular per acre. For use on lawns, apply to soil surface in form of a lawn fertilizer spreader. Treatment should be applied in the spring when insects are most likely to be troublesome. Do not treat a lawn if it is wet. Apply by broadcast or sprinkling.

CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water, and grass has completely dried. Birds and other wildlife in treated areas may be harmed.

USDA Reg. No. 476-1130

W WE 680606 SW

BURN EMPTY BAG — KEEP OUT OF SMOKE