UNDER THE FEDERAL INSECTIC HE FUNGICIDE AND RODENFICIDE AND RODENFICIDE TO ATTACHED COMMENTS.

OBSERVE THE FOLLOWING USE PRECAUTIONS

READ ENTIRE LABEL BEFORE USING.

Use only for crops listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed Federal tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Bo not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing of applications needed will vary with local conditions.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate any water supply or apply under conditions favoring drift outside the treated area.

 $^{\prime}$ This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

DIRECTIONS FOR USE

Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

Rinse spray equipment and containers. Bury liquid waste in pit in non-crop lands located away from water supplies.

DO NOT REUSE EMPTY CONTAINER. PERFORATE EMPTY METAL CONTAINERS. CRUSH AND BURY GLASS IN SAFE PLACE.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

1-ARC-7212J7-3

EPA Reg. No. 476-1142

LOT NO.

TOXAPH

AGRICULTURAL INSECTICIDE / EMULSIFIABLE LIQUID /

ACTIV	E IN	GREDI	E	STF	:
_				•	

Toxaphene (Technical Chlorinated Camphe	ene)*
Petroleum Hydrocarbon Solvent	
Xylene Range Aromatic Hydrocarbon Solve	
NERT INGREDIENTS:	

*Chlorine Content 67% to 69%.

CONTAINS 6 LBS. TOXAPHENE PER GALLI

WARNING

Keep Out of

MAY BE FATAL IF SWALLOWED.

MAY BE ABSORBED THROUGH SKIN.

DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON WASH AFTER HANDLING. DO NOT CONTAMINATE FEED OF

DO NOT USE, POUR, SPILE OR STORE NEAR HEAT OF PROTECT FROM TEMPERATURES BELOW 0°F. Product crystallizes at lot at higher temperatures and mix to re-dissolve crystals and assure unifor NOT FOR USE OR STORAGE IN OR AROUNI

MADE IN U.S.A. BY

STAUFFER CHEMICAL (
WESTPORT, CT 05880



GAL. NET

PHENE

ECTICIDE LIQUID

6-EH

ited Camphene)*it	
carbon Solvent	
	100.0%

INS 6 LBS. TOXAPHENE PER GALLON

Keep Out of Reach of Children

'ED. H skin.

ST. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING.

NOT CONTAMINATE FEED OR FOOD.

SPILL OR STORE NEAR HEAT OR OPEN FLAME.

LOW 6°F. Product crystallizes at lower temperatures. Warm or store ⊩dissolve crystals and assure uniformity before use.

TORAGE IN OR AROUND THE HOME.

MADE IN U.S.A. BY

CHEMICAL COMPANY STPORT, CT 06880

RECOMMENDATIONS

Dosage rates are given in terms of pints of Toxaphene 6-EH per acre in sufficient water to provide thorough coverage (usually 3-10 gals. water/acre by aircraft; 20-100 gals. water/acre by ground equipment). Except where otherwise shown, begin applications at the first sign of infestation and repeat as needed to maintain control but observe use limits given for crops. Maximum permissable rate per acre, expressed as TOXAPHENE 6-EH, is given in parenthesis, (), after each crop claim.

ALFALFA (Seed Crop): Leafhoppers, Lygus Bugs, Thrips Use 2 pts. per acre during bloom 3 to 4 weeks after bud treatment. Armyworms, Cutworms, Grasshoppers. Use 2 pts. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. (21/3 quarts)

CORN: Armyworms, Fall Armyworms. Use 2 pts. per acre when larvae are young. Cutworms. Apply 23-4 pts. Apply at base of plants and cultivate immediately after application. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain. (4 quarts)

SOYBEANS: Armyworms, Bean Leaf Beetles, Blister Beetle, Bollworm, Corn Earworm, Green Cloverworm, Velvetbean Caterpillar, use 3 to 51/3 pts. With aerial application, the desired volume of insecticidal concentrate should be diluted with water and applied at 3 to 4 gallons per acre. Do not apply within 21 days of harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry. (4 quarts)

SMALL GRAINS (Barley, Tats, Wheat, Rye): Armyworms, Fall Armyworms. Use 2 pts. Apply when larvae are young. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restrictions on use of grains. 121/3 quarts