

CAPTAN-TRITHION® 5-2 DUST

AN AGRICULTURAL FUNGICIDE-INSECTICIDE-ACARICIDE

ACTIVE INGREDIENTS

Captan: N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide	5%
Carbophenothion S-[[p-chlorophenyl]thio]methyl O,O-diethyl phosphorodithioate	2%
INERT INGREDIENTS:	93%
	100%

*U. S. Pat. Nos. 2,553,770 - 2,533,771 - 2,553,776. Not for use in export outside of the United States, its territories and possessions, Canada or Mexico.

**U. S. Pat. No. 2,793,224

CAUTION
Keep Out of Reach of Children

HAZARDOUS IF SWALLOWED

Avoid breathing dust, contact with skin and eyes.

Wear dust respirator and goggles.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

Do not contaminate feed and food.

Wash all contaminated clothing with soap and hot water before re-use.

ANTIDOTE—TRITHION, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. **Atropine is antidotal.**

OBSERVE THE FOLLOWING USE PRECAUTIONS

Toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Apply only to areas specified on the label.

Observe use limitations given on the label for specific crops.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damages by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

This product is to be applied only by experienced personnel using approved safety equipment. Apply when air is calm. To obtain most effective results, apply with a ground duster or by airplane. Apply at first sign of infestation and infection. Make thorough coverage of plants with a uniform deposit of dust.

For control of diseases persisting after the prescribed time limits have expired, an effective fungicide such as captan should be used alone, in a manner which will not leave residues in excess of established tolerances.

STRAWBERRIES

For control of *Botrytis* Fruit Rot (Gray Mold), Strawberry Aphid, Two-spotted Mites, apply 25-50 pounds per acre. Begin applications when new growth starts and repeat at 7 to 10 day intervals. Use higher dosage or shorten interval under conditions of frequent rains, heavy dew or other circumstances favorable to infection. Do not apply within 3 days of harvest.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Made in U.S.A.
Stauffer Chemical Company
New York

USDA Reg. No. 476-1278

DESTROY EMPTY CONTAINER — BURN — STAY OUT OF SMOKE

S-SF-670608-WC

