



# CAPTAN 80 SPRAY-DIP

476-1287  
ACCEPTED  
JUL 14 1970  
U.S. DEPARTMENT OF AGRICULTURE  
OFFICE OF PEST CONTROL AND PLANT QUARANTINE

**AGRICULTURAL FUNGICIDE FOR USE AS A DIP  
OR WASH ON FRUITS, VEGETABLES AND ORNAMENTALS.**

**ACTIVE INGREDIENT:**  
CAPTAN N-[[trichloromethyl]thio]-  
4-cyclohexene-1,2-dicarboximide 80%  
**INERT INGREDIENTS** ..... 20%  
100%

Not licensed for use in countries where valid patents still exist.

**CAUTION** Keep Out of  
Reach of Children

Avoid inhalation of dust or spray mist.  
Avoid prolonged or repeated contact with skin.  
Do not contaminate food and feed.

STORE IN COOL, DRY PLACE — PROTECT FROM EXCESSIVE HEAT.

Toxic to fish. Keep out of lakes, streams and ponds.

**BURN EMPTY BAG —  
KEEP OUT OF SMOKE**

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

### RECOMMENDATIONS AND DIRECTIONS FOR USE

Captan 80 Spray-Dip is a special microfine wettable powder for post-harvest use for the prevention of various rots and molds during storage of fruits and vegetables, and is also recommended as a preplant treatment for ornamentals.

**APPLES, APRICOTS, CANTALOUPE, CHERRIES, CUCUMBERS, GRAPEFRUIT, LEMONS, LIMES, MANGOES, NECTARINES, ONIONS (green and bulb), ORANGES, PEARS, POTATOES, TANGELOS AND TANGARINES:**

**Storage Rots and molds (Rhizopus, Botrytis, Gloeosporium).** Use 1½ lb. per 100 gal water. Apply as a spray or in dip tank.

When used as a dip, recharge wash solution periodically as volume in tank is depleted. When tank volume is reduced 25%, bring water back to volume and add ½ lb. Captan 80 Spray-Dip for every 25 gallons of water added. At end of every 8-hour shift, empty tank, flush, and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation in tank during dipping operations.

**AZALEA, CARNATION, CHRYSANTHEMUM:** Damping-off — Dip cuttings in a suspension of 10 teaspoonfuls per gal. of water before bedding.

**GLADIOLUS, TUBEROUS BEGONIA:** Damping-off, Rot — Use 5 tablespoonfuls per gal. of water. Dip bulbs for 30 minutes. Stir, drain, and plant.

LBS. NET

U.S.D.A. Reg. No. 476-1287

S SF-700216-WC

MADE IN U.S.A

**STAUFFER CHEMICAL COMPANY  
NEW YORK, NY 10017**

**LABEL SPECIMEN**