

# MANEB 8 DUST

## AN AGRICULTURAL FUNGICIDE

### ACTIVE INGREDIENT:

Maneb (Manganese Ethylene Bisdithio carbamate) . . . . . 8%

### INERT INGREDIENTS: . . . . .

92%

100%

ACCEPTED

APR 24 1969

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 496-1333

### DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops.

### RECOMMENDATIONS

This dust is formulated for the control of the following diseases: Apply 25 to 35 lb. per acre, unless otherwise specified.

CARROTS: Leaf Blight, Cercospora Leaf Spot --- Start applications when plants are six weeks old (or when disease threatens) and continue every 7 to 10 days.

CELERY: Early and Late Blight --- Start applications when plants are established in the field and continue as frequently as required. CAUTION: Remove residues by stripping, trimming and washing.

CUCURBITS: (Cantaloupes, Cucumbers, Squash) Anthracnose and Downy Mildew. Start applications when runners begin to form or at the first sign of the disease. Do not apply unless plants are dry.

ONIONS: Botrytis Leaf Blight, Downy Mildew, Purple Blotch --- Start application when plants are six weeks old (or when disease threatens) and continue at 7 to 10 day intervals as long as necessary.

POTATOES: Early and Late Blight --- Start application as soon as the plants are 2 to 6 inches high and continue at 5 to 10 day intervals as long as plants threaten.

SPINACH: Downy Mildew (Bluemold) --- Apply 25 to 35 lb. per acre at the first sign of mildew and repeat at 7 to 10 day intervals or as needed for adequate control. Do not apply within 7 days of harvest. Remove residues by washing.

TOMATOES: Early and Late Blights, Anthracnose, Gray Leaf Spot, Septoria Leaf Spot --- Start application when fruit or fruit clusters are 1/2 to 1 inch and repeat at regular intervals.

### CAUTION

#### Keep Out of Reach of Children

- May cause irritation of nose and throat.
- Do not breath dust.
- Wear respirator.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

This product is toxic to fish. Keep out of lakes, streams and ponds.

Consult state agricultural extension service or state agricultural experiment stations for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.