

TRITHION-SULFUR 2-50 DUST

ACCEPTED
12-1-69
476-1504

ORGANIC PHOSPHATE INSECTICIDE-ACARICIDE & FUNGICIDE

ACTIVE INGREDIENTS:

Carbophenothion—S-[[p-chlorophenyl]thio]methyl
O,O-diethyl phosphorodithioate
Sulfur

2%
50%
48%
100%

INERT INGREDIENTS:

U. S. Pat. No. 2,793,224

Reg. Stauffer Chemical Co. for insecticide-acaricide.

CAUTION

Keep Out of Reach of Children

HAZARDOUS IF SWALLOWED. AVOID BREATHING DUST, CONTACT WITH SKIN AND EYES. Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate food and feed. Wash all contaminated clothing with soap and hot water before re-use.

TRITHION, upon repeated, prolonged or careless use may cause cholinesterase inhibition. **ATROPINE IS ANTIDOTAL.** Keep supply of Atropine 1/100 grain tablets on hand.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Apply only on recommended crops.

Do not apply sulfur when temperature is high as injury to foliage may result.

Avoid dust drift to adjoining food and forage crops.

Toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Apply only to areas specified on the label.

Apply in early morning or evening when there is little or no wind.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service, as the timing, number, and rate of applications needed will vary with local conditions.

BURN EMPTY BAG—KEEP OUT OF SMOKE

DIRECTIONS FOR USE

This product is to be applied only by experienced personnel using approved safety equipment. Apply at first sign of infestation and infection by ground equipment or aircraft. Unless otherwise stated repeat at 7 to 10-day intervals as needed but observe use limitations given for specific crops.

For control of diseases persisting after the prescribed time limits have expired, an effective fungicide should be used alone, in a manner which will not leave residues in excess of established tolerances.

RECOMMENDATIONS

ALFALFA: Two-spotted mite, pea aphid, spotted alfalfa aphid, potato leafhopper, meadow spittlebug—Apply 25 to 50 lbs per acre. Use 50 lbs per acre for the control of the meadow spittlebug. Do not apply more than one application per cutting. Do not graze or cut hay within 28 days after application.

BEANS, DRY (shelled): Bean aphid, Mexican bean beetle, potato leafhopper, powdery mildew, two-spotted mite, serpentine leaf miner—Apply 30 lbs per acre. Do not apply more than two applications per season. Do not graze or cut hay within 21 days after application.

BEAN, SNAP: Bean aphid, serpentine leaf miner, potato leafhopper, Mexican bean beetle, powdery mildew, two-spotted mite—Apply 30 lbs per acre. Do not apply within 7 days of harvest. Do not apply more than two applications per season. Do not graze or cut hay within 21 days after application.

CHERRY, PEACH, PLUM, PRUNE: Aphids, orchard mites, powdery mildew—Apply 40 to 50 lbs per acre. Do not apply within 30 days of harvest.

CLOVER: Two-spotted mite, pea aphid, potato leafhopper, meadow spittlebug—Apply 25 to 50 lbs per acre. Use 50 lbs per acre for the control of the meadow spittlebug. Do not apply more than one application per cutting. Do not graze or cut hay within 28 days after application.

GRAPES: Grape aphid, grape leafhopper, grape vine mealybug—Apply 40 to 50 lbs per acre. Do not apply within 30 days of harvest. Do NOT APPLY TO SENSITIVE

VARIETIES SUCH AS: CHASSARD, CARMY, GOLDEN MUSCAT, MUSCA, AND ITALIA.

PEA: Pea aphid, powdery mildew, serpentine leaf miner, two-spotted mite—Apply 30 to 40 lbs per acre. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

POWDERY MILDWEE ON CROPS (including alfalfa): Apply 40 to 50 lbs per acre. Do not apply within 30 days of harvest. Do not graze or feed crop residue to livestock.

SPINACH: Spinach aphid, two-spotted mite—Apply 30 to 40 lbs per acre. Do not apply within 14 days of harvest. Crop residue can be fed to livestock.

STRAWBERRY: Powdery mildew, strawberry aphid, two-spotted mite—Apply 30 to 40 lbs per acre. Do not apply within 7 days of harvest.

SUGAR BEETS: Pea aphid, potato flea beetle, serpentine leaf miner, two-spotted mite—Apply 30 to 50 lbs per acre. Do not apply within 14 days of harvest. Tops are to be used for feed.

TOMATO: Potato flea beetle, potato psyllid, potato leafhopper, tomato russet mite, tropical mite, serpentine leaf miner, two-spotted mite—Apply 25 to 50 lbs per acre. Do not apply within 7 days of harvest.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

50 LB. NET

Made in U.S.A.
Stauffer Chemical Company
New York

USDA Reg. No. 476-1504
S-SF-681216-NP