# GAPTANIESBEMANIES 20-20-20 SEED DAOTEGIANI

#### FOR SEED DISEASE CONTROL

#### SPECIAL LUBRICANT ADDED - DISCOLORED

## DRILL BOX OR PLANTER BOX APPLICATION

Captan: N-[(Trichloromethyl)thio]-4-cyclohexene- 1,2-dicarboximide*								20%
1,2-uicarboximide	•	•	•	•	•	•	•	200/
Hexachlorobenzene	•	•	•	•	•	•	•	20%
Maneb: Manganese ethylenebisdithiocarbamate	•		•	•			•	20%
INERT INGREDIENTS	•	•	•	•	•	•	-	40 /0
								100%

### CAUTION

Keep Out of Reach of Children

#### MAY CAUSE IRRITATION OF NOSE, THROAT, EYES AND SKIN.

Do not breathe dust. Wear respirator, goggles and gloves.

Avoid confact with skin, eyes and clothing. In case of contact with eyes, flush with plenty of water and get medical attention.

Wash thoroughly after using.

Avoid contamination of food or feed.

Do not use treated seeds for human consumption or for livestock feed.

Toxic to fish. Keep out of lakes, streams or ponds, do not contaminate by cleaning equipment, or disposal of wastes.

#### USE PRECAUTIONS

Exact all directions before using and use only as specified on this label. If an container tightly closed when not in use. Keep away from fire or sparks. Stare in a cool, dry place. Do not allow to become wet or over-heated in the cool.

has may bring on chemical change which will impair the fungicidal effectiveness of the product and may also produce flammable vapors.

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES.

NCTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

#### PRODUCT INFORMATION

Use only recommended rates. Lower amounts may negree desired control. Excessive amounts may cause seed injury. This material is a special seed treatment formulation of Captan, HCB and Maneb which, when used at the correct application rates, protects seeds and seedlings against seed and soil-borne diseases such as seed decay, root rot, damping-off and seedling blight. Also, protection is given against stinking smut of wheat, oats smut, covered and false loose smuts of barely and bunt of rye.

#### RECOMMENDATIONS and DIRECTIONS FOR USE

#### PLANTER BOX OR DRILL BOX APPLICATION RATES

Use on seed by mixing with the seed in the box at the rate shown below. DO NOT MIX WITH HANDS.

Wheat, oats, barley, rye	3 to 0 oz. bu.
Corn seed (field)	3 to 0 62. ba.
Soybeans, beans, peas	3 to 3 oz. bu.
Peanuts (shelled)	3 to 6 oz. inu.
Cotton	3 to 6 uz. 10u#

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled.

## TREATED SEED DO NOT USE FOR FOOD, FEED OR OIL PURPOSES

The federal government and some states have issued regulations to the application of special tags or labels to be affixed to bags containing treated seed.

USDA Reg. No. 4/6-2082

ARC-710311-MW

FOR ECOMONIC + .. SQU REGISTER.

SO UNDER NOT A LACABOR

50 LB. NET

STAUFFER CHEMICAL COMPANY