

CAPTAN-SEVIN*

7.5-2.5 DUST

AGRICULTURAL INSECTICIDE AND FUNGICIDE

ACTIVE INGREDIENTS:

Captan: *N*-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide** 7.5%
Carbaryl: (1-naphthyl *N*-methylcarbamate) 2.5%

INERT INGREDIENTS:

. 90.0%
. 100.0%

*Registered trademark of Union Carbide Corporation for active ingredient (carbaryl).

**Not licensed for use in countries where valid patents still exist.

CAUTION

Keep Out of Reach of Children

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing dust. Avoid contact with skin. Wash hands and face before eating. Shower or bathe after work. Wear long sleeved work clothing. Change to clean clothing daily. Do not contaminate food or feed.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

STORE IN A COOL DRY PLACE. PROTECT FROM EXCESSIVE HEAT.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

BURY WASTE AND EMPTY CONTAINER IN A SAFE PLACE

50 LBS. NET

EPA REG. NO. 476-2113-AA

ARC-720313-0

OBSERVE THE FOLLOWING USE PRECAUTIONS

Read all label directions before using. Use only for claim listed and only as specified on this label.

This product is highly toxic to humans exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

SEVIN insecticide will not control insects if used with lime or other highly alkaline materials.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crop as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with the local conditions.

FISH CAUTIONS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7 to 10 day intervals when needed to maintain control.

RECOMMENDATIONS

STRAWBERRIES: Botrytis Fruit Rot (Gray Mold), Leaf Spot, Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil - apply 40 to 50 pounds per acre. May be applied up to 1 day before harvest.

Made In U.S.A. By
STAUFFER CHEMICAL COMPANY
Westport, Connecticut 06880