

199 68 ED  
 EQUIPMENT UNDER FEDERAL INSPECTION  
 TUBERCULOCIDAL  
 FICIDE ACT  
 COMMENTS REGISTER

**STAPHYLOCIDAL  
 TUBERCULOCIDAL**

STAPH-KILL will control the following organisms: Tubercle bacilli, Salmonella typhosa, Shigella paradysenteriae, Salmonella schottmulleri, Escherichia coli, Staphylococcus aureus, Streptococcus pyogenes, Mycobacterium smegmatis, Trichophyton interdigitale (ringworm or athlete's foot), Penicillium glaucum, Influenza A<sub>2</sub>, and Mumps Viruses.

STAPH-KILL is a powerful hospital disinfectant and deodorant for germ laden surfaces, . . . Metal Beds and Springs, Wheel Chairs, Linen Carts, Hampers, Refuse Cans, Dressing Carts, Cleanup Carts, Unpolished Furniture, Air Ducts, Toilet Seats, Telephone, Wash Basins, Urinals, Light Switches, Office Machinery, Laundry Chutes, Door Knobs, Athletic Equipment, Language Laboratory Equipment. . . .

STAPH-KILL will reduce the hazard of cross infection by bacteria and viruses due to contaminated surfaces.

STAPH-KILL is a QUICK ACTING, non-selective odor killer which acts by chemically neutralizing odor producing materials. Excellent for sanitizing and deodorizing patients' rooms, class rooms, meeting rooms, locker rooms and other areas.

USDA REG. No. 491-199



**NEW  
 FUNGICIDAL  
 BACTERICIDAL**



A Division of National Service Industries, Inc.

Atlanta	Miami	Louisville
Dallas	New Orleans	Los Angeles
Houston		San Juan

**CAUTION:**  
 KEEP OUT OF REACH OF CHILDREN.  
 Contents under pressure. See side panel for additional required precautions.

Alcohol Registration No. SDA 11-235

NET WGT. 15½ OZ. AV.

USDA REG. No. 491-199

**CAUTION:  
 FLAMMABLE**

Contents under pressure. Avoid contact with food or foodstuffs. Store away from heat. Do not puncture or dispose of empty container in fire. Do not use on polished wood, furniture, or rayon fabrics.

**SHAKE WELL  
 BEFORE USING**

For Disinfecting: After cleaning object, spray surface until wet from a distance of 6 to 8 inches.

For Deodorizing: Hold dispenser upright; point valve opening away from face. Spray into air towards center of room for 3 to 4 seconds.

ACTIVE INGREDIENTS:  
 Orthophenylphenol .08%  
 Para Tertiary Amyl Phenol . . . .02%  
 Ethyl Alcohol . . . .62.40%  
 INERT INGREDIENTS . 37.50%