


Scotts

house plant insect spray



Controls major damaging insects and protects house plants

Advantages

- **Knocks down** common house plant insects and protects plants from further damage up to two weeks.
- **For use on flowering and foliage house plants** anytime of the year.
- **Safe for plants** when used as directed.
- **Convenient,** ready to use spray bottle.
- **Research tested by Scotts.**
- **Guaranteed—Scotts No-Quibble Guarantee.**
- **For more information,** call or write.

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUTIONARY STATEMENTS

See other side for additional cautions.

NET WT. 8 FL. OZ. (24L)

ACTIVE INGREDIENT:

cyano (3-phenoxyphenyl) methyl
4-chloroalpha (1-methylethyl)
benzeneacetate —01%
INERT INGREDIENTS 99.99%
Total 100.00%

EPA Reg No 538-166 EPA Est No 538-OH-1
Product of USA
The O M Scott & Sons Company
Marysville, OH 43041

Hazard to Humans and Domestic Animals: Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain

medical attention if irritation persists. Do not apply to plants intended for consumption. Do not mix with other pesticides. Do not apply to food. Remove pets, birds, and cover fish aquariums before spraying.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Apply this product only as specified on the label.

Storage and Disposal: Do not contaminate water, food or feed by storage or disposal.
Storage: Store in a cool dry place. Avoid freezing or excessive temperatures.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

Scotts house plant insect spray

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

When: Anytime of year when insects are observed.

Where: To all leaf surfaces, top and bottom, including new, emerging leaves, stems, and the junction of stems and leaves. Try to contact the insects with the spray. Avoid spraying blossoms of flowering plants since spray may discolor blossoms.

How Much: Shake first, then hold container 10-12 inches from plant and spray a fine mist. Stop just before the spray begins to run off. Repeat in 14 days. For heavy infestation, reapply in 7 days.

SUGGESTIONS

• For improved control, remove as many insects as possible with a swab before spraying.

- For prevention on newly acquired plants, spray twice, 7 days apart, to control insects that may be present and thus avoid exposing insect-free plants in the home.

- When practical and when temperatures are above 55°, take plants outside for spraying or into a separate area; e.g. a patio or garage, for spraying. Avoid spraying plants while they are in direct sunlight or if they are wet from misting.


Scotts Policy No-Quibble Guarantee. If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase, and we will mail you a refund check promptly.

Scotts . . . Marysville, Ohio 43041

©1982 Distributed by The O M Scott & Sons Company
All rights reserved. XXXX

820523

Printed in U.S.A.



181