

NET WEIGHT 50 LBS.

BOTRAN® 6 DUST

FUNGICIDE

ACTIVE INGREDIENT:

2,6-Dichloro-4-Nitroaniline

INERT INGREDIENTS:

Total

6%

94%

100%

U.S.D.A. Reg. No. 550-42

WARNING! HARMFUL IF SWALLOWED. KEEP OUT OF REACH OF CHILDREN.

Avoid inhalation of dust. In case of contact, remove contaminated clothing and wash skin with soap and water. PRECAUTION: This product toxic to fish, keep out of any body of water. Do not use on germinating seeds or annual seedlings except as suggested. Avoid contamination of

feed and foodstuffs.

Do not use this product for any purpose other than that recommended on this label. Do not allow this product to drift from the area being treated.

Recommendations for Use:

APRICOT: Use 50 lbs. per acre to control Fruit Decay (Brown Rot *Monilinia* and *Rhizopus* Rot).

Apply 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest.

LETTUCE: (Head) Use 30 lbs. per acre to control Wilt (*Botrytis* Rot and *Sclerotinia*). Apply first application at thinning. Repeat application 7 days later. Do not apply within 14 days of harvest.

PEACHES AND NECTARINES: Use 60 lbs. per acre to control Blossom Blight *Monilinia* and Fruit Decay (Brown Rot *Monilinia* and *Rhizopus* Rot). For Blossom Blight, apply at pink bud and full bloom. For Fruit Decay, apply at 18 days, 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest.

SWEET CHERRY: Use 50 lbs. per acre to control Blossom Blight *Monilinia* and Fruit Decay (Brown Rot *Monilinia* and *Rhizopus* Rot). For Blossom Blight, apply at popcorn, bicora, full bloom and petal fall. For Fruit Decay, apply 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest.

CANEBERRIES: (Blackberries, Boysenberries, Red Raspberries) Use 50 lbs. per acre to control Fruit Decay (*Botrytis* Rot). Begin applications at first appearance of symptoms. Repeat at 10 day intervals. Use a maximum of 4 applications. Do not apply within 7 days of harvest.

ONION AND GARLIC: Use 250-400 lbs. per acre to control White Rot (*Sclerotinia*). Direct seeding -- incorporate Botran in top 1 1/2 inches of soil 1-2 weeks before onions are seeded. Planting sets or buds -- dust Botran into furrow when planting so treated soil surrounds sets or buds. Rate per acre will vary in local areas. Check local agricultural authorities for more exact recommendation.

ROSE AND HYDRANGEA: Use 30 lbs. per acre to control Blight (*Botrytis*). Field -- Apply Botran prior to shipment or storage. Stored -- Thoroughly dust storage area and all canes and dead foliage; repeat application during long storage periods.

GLADIOLUS: Use 250 lbs. per acre to control Dry Rot (*Strombotrytis*). Dust into furrow at planting time, so treated soil surrounds corms. Use 30 lbs. per acre to control Blight (*Botrytis*). Dust thoroughly. Force fungicide between spikes. Apply to bunched cut flower spikes after harvest and before shipment or storage.

ACCEPTED

FEB 27 1970

UNDER THE FEDERAL FUNGICIDE ACT FOR ECONOMIC REGULATION UNDER NO. 550-42 TO ATTACHED COMMENT.

SELLER MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL OR THE USE OF THIS PRODUCT OTHER THAN INDICATED ON THE LABEL. BUYER ASSUMES ALL RISK OF USE AND/OR HANDLING OF THIS MATERIAL WHEN SUCH USE AND/OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS.



Van Waters & Rogers

550 42