



BOTRAN[®] 50% DUST FUNGICIDE

ACTIVE INGREDIENTS	2,6-Dichloro-4-Nitroaniline	50%
INERT INGREDIENTS:	50%
	Total	100%

U.S.D.A. Reg. No. 550-86

BOTRAN[®] Reg. T.M. Tuco Products Co., Div. of the Upjohn Co.

WARNING! KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED.
Avoid inhalation of dust. In case of contact, remove contaminated clothing and wash skin with soap and water.

PRECAUTION: This product toxic to fish, keep out of any body of water. Not for use in the North Central States.

DIRECTION FOR USE

ONION AND GARLIC: Use 30 - 48 lbs. per acre or per 43,560 linear feet of row to control white rot (Sclerotium). Direct seeding — Incorporate Botran in top 1½ inches of soil, 1 to 2 weeks before seeding onions. Planting Sets or Cloves — Dust Botran into furrow when planting so created soil surrounds sets or cloves. Rate per acre will vary in local areas. Plow soil to 8 inches and cross disk the area prior to seeding sensitive follow-up crops.

Do not plant spinach as a follow-up crop in treated soil.
Do not use on germinating seeds or annual seedlings except as suggested.
Do not use this product for any purpose other than that recommended on this label. Do not allow this product to drift from the area being treated. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Distributed by



Van Waters & Rogers

SAN FRANCISCO, CALIFORNIA 94119

NET WEIGHT 50 POUNDS

SELLER MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL OR THE USE OF THIS PRODUCT OTHER THAN INDICATED ON THE LABEL. BUYER ASSUMES ALL RISK OF USE AND/OR HANDLING OF THIS MATERIAL WHEN SUCH USE AND/OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS.

Printed in U.S.A. (5/70)

