

CONCENTRATE

FOR PEST CONTROL OPERATORS ONLY

ACTIVE INGREDIENTS:

Pyrethrins	1.0%
Technical Piperonyl Butoxide*	10.0%
Petroleum Hydrocarbons	89.0%

*Equivalent to 9.0% of (butylcarbityl (di-propyl) piperonyl) ether and 2.0% related compounds

EPA Reg. No. 550-125-AA

EPA Est. No. 5316-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Avoid breathing spray mist. Wear a mask or respirator of a type recommended by NIOSH to give adequate protection against this material. Wash hands, arms, and face with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Remove persons and pets prior to use and ventilate treated areas prior to re-entry. Remove or cover fish bowls (tanks) before spraying or fogging. Do not make applications when food is present or unprotected. Food handling utensils and equipment should be well protected or washed thoroughly before use.

COMBUSTIBLE MIXTURE: Do not use, pour, spill, or store near heat or open flame. Keep container closed.

CONTAINER DISPOSAL: Do not reuse empty container. Dispose of in accordance with state and local laws and regulations. Where permitted, destroy this container by perforating or crushing and then burying or discarding in a safe place away from water supplies. Empty drums may be sent to an approved reconditioner.

FISH & WILDLIFE: This product is poisonous to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on labeling.

Apply, use, store, handle only as specified on product labeling. Buyer/User may be subject to civil and/or criminal penalties for failure to observe the labeling directions and precautions.

NET CONTENTS: 1 GALLON

Manufactured By

NAMCO

Van Waters & Rogers

765 Landess Avenue, Milpitas, CA 95035

Apply only as specified on labeling.

DIRECTIONS FOR USE

For use in homes, apartments, offices, dairies, hospitals, warehouses, restaurants, and food processing plants

For maximum effectiveness, a combination of localized surface application and space treatment is recommended. Repeat treatments as necessary.

When treating for stored product pests remove and destroy all infested materials, where possible. Fumigate, or treat by other effective means, using an approved product intended for this purpose for control of non-exposed or accessible insects.

AS A SURFACE SPRAY — For initial clean-up of severe insect infestations, dilute at a rate of 1 part concentrate in 4 parts water or suitable spray base oil. This dilution may not be used in edible product areas of federally inspected meat and poultry plants. In these plants and for normal infestations, dilute 1 part concentrate in 9 parts water or oil.

AS AN AID IN THE CONTROL OF EXPOSED OR ACCESSIBLE STAGES OF THE FOLLOWING HOUSEHOLD AND STORED PRODUCTS PESTS: COCKROACHES, ANTS, SILVERFISH SPIDERS, CRICKETS, BOX ELDER BUGS, CLOVER MITES, WEEVILS (GRANARY, BEAN, RICE), BEETLES (CONFUSED AND RUSTY RED FLOUR, CIGARETTE, DRUGSTORE, FLAT, SAW-TOOTHED AND RUSTY GRAIN), LESSER GRAIN BORER, MEAL WORMS, GRAIN MITES, CADELLES, MOTHS (MEDITERRANEAN FLOUR, INDIAN MEAL, ANGOUMIS GRAIN). Adjust spray nozzle to give a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under panels, around containers of stored foods, around the bases of machinery, behind shelves and drawers. For ants, spray trails, nests and point of entry. All surface application only. To be used on spray treated walls and other surfaces applying at the rate of 1 quart per 100 square feet of area.

TO CONTROL FLEAS — Thoroughly spray infested area (pet beds, resting quarters, kennels, crates and previous latrine and bedding quarters) and treat with low temperature steam and utilize fumigation of food and feed rations. For flea doghouses, 1 to 2 pints per gallon of water for treatment. Only repeat treatment if necessary by reapplying to areas not treated. Do not treat animals with this formulation.

AS A SPACE SPRAY — AS AN AID IN THE CONTROL OF EXPOSED OR ACCESSIBLE STAGES OF FLIES, FRUIT FLIES, GNATS, MOSQUITOS, SKIPPER FLIES, WASPS, HORNETS, MOTHS (MEDITERRANEAN FLOUR, INDIAN MEAL, ANGOUMIS GRAIN). This product may be applied undiluted or diluted to the maximum rates shown below.

Conventional spray equipment — 1 to 20

Vaporizing equipment (non-thermal) — 1 to 10

Fogging equipment (thermal) — 1 to 5

Dilute with suitable spray base oil only.

USE PRECAUTIONS

TREAT FOOD PROCESSING PLANTS ONLY WHEN PLANTS ARE NOT IN OPERATION. When used in meat and poultry processing plants, this product must be diluted at least 1 to 1 with a suitable spray base oil. Consult table above for maximum dilution rates. The spray mixture must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

Prior to Treatment

Remove or cover exposed food and food handling surfaces.

Remove all persons, birds and other pets from the area to be treated. Cover fish bowls (tanks) if they cannot be removed.

Close all doors and windows. Shut off ventilation system, air ducts and all other air conditioning equipment.

Enter the plant facility wearing protective

clothing. Apply the concentrate to the target area. Avoid contact with the concentrate. Do not breathe the spray. Do not ingest or drink. Do not eat or drink. Do not use tobacco products.

Do not use in areas where food is prepared.

Do not use in areas where food is stored.

After application, ventilate the area. Remove all persons, birds and other pets from the area. Remove all protective clothing and wash thoroughly with soap and water. Do not use tobacco products.

Do not use in areas where food is prepared. Do not use in areas where food is stored. Do not use in areas where food is handled. Do not use in areas where food is displayed.

For more information, contact NAMCO.

NAMCO is a trademark of NAMCO Van Waters & Rogers