

Proposed labeling w/ USDA-APHIS approval
 PM 17 RIG# 550-125



NAMCO®

PERMETHRIN FOG CONCENTRATE

FOR PEST CONTROL OPERATORS ONLY

ACTIVE INGREDIENTS:

Permethrin	1.0%
Technical Piperonyl Butoxide*	10.0%
Petroleum Hydrocarbons	89.0%

*Equivalent to 8.0% of (butylcarbityl) (6-propylpiperonyl) ether and 2.0% related compounds

EPA Reg. No. 550-125-4A
 EPA Est. No. 5316-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Avoid breathing spray mist. Wear a mask or respirator of a type recommended by MSDS to give adequate protection against this material. Wash hands, arms, and face with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Remove persons and pets prior to use and ventilate treated areas prior to re-entry. Remove or cover fish bowls (tanks) before spraying or fogging. Do not make applications when food is present or unprotected. Food handling utensils and equipment should be well protected or washed thoroughly before use.

COMBUSTIBLE MIXTURE: Do not use, pour, spill, or store near heat or open flame. Keep container closed.

CONTAINER DISPOSAL: Do not reuse empty container. Dispose of in accordance with state and local laws and regulations. Where permitted, destroy this container by perforating or crushing and then burying or discarding in a safe place away from water supplies. Empty drums may be sent to an approved recycler.

FISH & WILDLIFE: This product is poisonous to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on labeling.

Apply, use, store, handle only as specified on product labeling. Buyer/User may be subject to civil and/or criminal penalties for failure to observe the labeling directions and precautions.

NET CONTENTS: 1 GALLON

Manufactured by

NAMCO®
Van Waters & Rogers

706 London Avenue, Milpitas, CA 95035

Apply only as specified on labeling.

DIRECTIONS FOR USE

For use in homes, apartments, offices, dairies, hospitals, warehouses, restaurants, and food processing plants.

For maximum effectiveness, a combination of localized surface application and space treatment is recommended. Repeat treatments as necessary.

When treating for stored product pests remove and destroy all infested materials, where possible. Fumigate, or treat by other effective means, using an approved product intended for this purpose for control of non-exposed or accessible insects.

AS A SURFACE SPRAY — For initial clean-up of severe insect infestations, dilute at a rate of 1 part concentrate in 4 parts water or suitable spray base oil. This dilution may not be used in edible product areas of federally inspected meat and poultry plants. In these plants and for normal infestations dilute 1 part concentrate in 9 parts water or oil.

AS AN AID IN THE CONTROL OF EXPOSED OR ACCESSIBLE STAGES OF THE FOLLOWING HOUSEHOLD AND STORED PRODUCTS PESTS: COCKROACHES, ANTS, SILVERFISH, SPIDERS, CRICKETS, BOX ELDER BUGS, CLOVER MITES, WEEVILS (GRANARY, BEAN, RICE), BEETLES (CONFUSED AND RUSTY RED FLOUR, CIGARETTE, DRUG-STORE, FLAT, SAW-TOOTHED AND RUSTY GRAIN), LESSER GRAIN BORER, MEAL WORMS, GRAIN MITES, CADELLES, MOTHS (MEDITERRANEAN FLOUR, INDIAN MEAL, ANGOUMIS GRAIN). Adjust spray nozzle to give a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored goods and around the bases of machinery behind shelves and drawers. For ants, spray trails, nests and point of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at the rate of 1 gallon to 750 square feet of surface.

TO CONTROL FLEAS: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor coverings. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals by a veterinarian is recommended. Do not treat animals with this formulation.

(continued on side/reverse panel)

SELLER MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL OR THE USE OF THIS PRODUCT OTHER THAN INDICATED ON THE LABEL. BUYER ASSUMES ALL RISK OF USE AND/OR HANDLING OF THIS MATERIAL WHEN SUCH USE/OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS.

©1976 NAMCO, a division of VAN WATERS & ROGERS

ACCEPTED
 Oct 7 1976
 United States Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 550-125

AS A SPACE SPRAY — AS AN AID IN THE CONTROL OF EXPOSED OR ACCESSIBLE STAGES OF FLIES, FRUIT FLY, GNATS, MOSQUITOS, SKIPPER FLIES, WASPS, HORNET, MOTHS (MEDITERRANEAN FLOUR, INDIAN MEAL, ANGOUMIS GRAIN). This product may be applied undiluted to the maximum rates shown below:
 Conventional spray equipment — 1 to 20
 Vaporizing equipment (non-thermal) — 1 to 10
 Fogging equipment (thermal) — 1 to 5
 Dilute with suitable spray base oil only.

USE PRECAUTIONS

TREAT FOOD PROCESSING PLANTS ONLY WHEN PLANT ARE NOT IN OPERATION. When used in meat and poultry processing plants, this product must be diluted at least 1 to 10 with a suitable spray base oil. Consult label above for maximum dilution rates. The spray mixture must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 5 microns in diameter and 80% less than 30 microns).

Prior to Treatment


- Remove or cover exposed food and food handling surfaces.
- Remove all persons, birds and other pets from the area to be treated. Cover fish bowls (tanks) if they cannot be removed.
- Close all doors and windows. Shut off ventilation system, air ducts, and air conditioning equipment.
- Extinguish all open flames including pilot lights.

During Application

- Avoid breathing spray mixture. Use a respirator.
- Fill room completely with the mist or fog by directing toward the upper portions of the room and corners.

After Application

- Do not remain in treated area.
- Leave room closed for at least 10 minutes.
- Ventilate thoroughly prior to re-entry in hospital and/or processing areas.
- Wipe up all exposed food processing surfaces and wash thoroughly before use.



OLD
[5316-43]

Apply only as specified on this label.

DIRECTIONS

SPACE SPRAY: For control of flies, mosquitoes and gnats. See use precautions below. Mix 4 1/2 fluid ounces of the 3.0% emulsifiable concentrate in a gallon of water to make a 0.10% emulsion. Mix well before using. Apply as a fine spray at the rate of 2 fluid ozs. per 1,000 cu. ft. or 1 gallon per 64,000 cu. ft.

USE PRECAUTIONS

TREAT FOOD PROCESSING PLANTS ONLY WHEN PLANTS ARE NOT IN OPERATION.

Prior to Treatment:

Remove or cover exposed food and food handling surfaces.

Remove all persons, fish, birds and other pets from the area to be treated.

Close all doors and windows. Shut off ventilation system, air ducts, and air conditioning equipment.

Extinguish all open flames including pilot lights.

During Application:

Avoid breathing mist. Use a respirator.

Fill room completely with the mist by directing toward the upper portions of the room and corners.

To reduce staining, spray exit should be a minimum of 5 feet from objects.

After Application:

Do not remain in treated area.

Leave room closed for 30 minutes.

Ventilate thoroughly prior to re entry. In hospitals, ventilate the room for a minimum of 2 hours.

Wash all exposed food processing surfaces and utensils thoroughly before use.

Repeat as necessary.

CONTACT SPRAY: For control of roaches, flies, mosquitoes and gnats. Mix 1/2 cup of

For use
hotels,
handling

ACTIVE

• (5 Be

2 met

Relate

Aromat

INERT

NE

Fresh