#### DIRECTIONS FOR USE

THE CONTRACT OF THE PROPERTY OF THE PARTY OF

General: This product is intended for use in control of brush and woody plants, and should not be used for weed control in growing crops.

Foliage Spray: For woody growth less than 6 feet tall, dilute at rate of 3 quarts per 100 gallons of water and spray bark and foliage thoroughly. Spray when plant is in active growing condition. If new growth appears, repeat application.

**Stump Treatment:** For plants 6 feet or higher, cut near ground. Dilute at rate of 3 to 4 quarts per 25 gallons of fuel oil or kerosene and spray bark and cut surfaces.

**Dormant Treatment:** Dilute as for stump treatment and spray brush stems and trunks thoroughly to height of 12 to 18 inches.

Mesquite: Dilute 1 gallon in 5 gallons of diesel fuel and emulsify with 18 gallons of water. Apply mixture at rate of 4 gallons per acre when plants have reached first heavy foliage stage of growth in spring (40-90 days after first leaves appear).

NET CONTENTS

# 4T Amine Brush Killer

Contains 4 lbs. 2, 4, 5-T Acid Equivalent Per Gallon

 ACTIVE INGREDIENTS:

 TRIETHANOLAMINE 2, 4, 5-TRICHLO 65.0%

 ROPHENOXYACETATE (Equivalent to 41% of 2, 4, 5-T Acid)
 65.0%

 INERT INGREDIENTS (100.0%)
 35.0%

For control of many kinds of brush and woody plants such as Alder, Aspen, Birch, Chokeberry, Elder, Elderberry, Elm, Hickory, Locust, Poison Ivy, Poison Oak, Sumac, Wild Cherry, Wild Grape and Willow, on range lands, rights-of-way, stump land and waste land areas where it is desired to kill or repress growth of woody plants.

### **CAUTION**

#### KEEP OUT OF REACH OF CHILDREN

#### (SEE BACK OR SIDE PANEL FOR OTHER PRECAUTIONS)

Do not take internally. May cause skin irritation. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Because of difficulty of cleaning sprayers and other equipment used it is recommended that such equipment not be used for handling and applying other agricultural chemicals. Do not store near fertilizers, seeds, insecticides or fungicides.

Vapors from this product may injure plants in the immediate vicinity.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

USDA REG. NO. 551-116



Manufactured by
BAIRD & McGUIRE, INC.
HOLBROOK, MASSACHUSETTS

## ACCEPTED

and the state of t

SEP 28 1970

UNDER THE TOTAL INSECTIONS
FUNGICIDE AT A RODENTICIDE ACT
FOR ECONO ATO POISON, REGISTER.
ED UNDER NO. TO SUBJECT
TO ATTACHED COMMENTS.

#### OBSERVE THESE PRECAUTIONS

- Do not apply directly to, or otherwise permit contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other desirable plants.
- 2. Avoid spray drift as even minute quantities of the spray may cause severe injury during both growing and dormant periods. (Coarse sprays are less likely to drift.)
- 3. Do not use on lawns or pastores entitioned has
- 4. Do not contaminate irrigation water or w. ter used for domestic purposes. Excessive amounts of 4T-Amine Brush Killer in the soil may temporarily inhibit seed germination and plant growth.
- Avoid storing near fertilizers, seeds, insecticides or fungicides.
- 6. Do not reuse this container for any other material
- 7. Do not reuse spray rig which has been used for spraying or storing of 2, 4, 5-T solutions.
- Use wisely and according to directions. Consult your local agricultural experiment station for specific applications.

Note ONLY NAME

. .

. .. .

# 17 Amine Brush Killer

Contains 4 lbs. 2, 4, 5-1 Acid Equivalent Per Gallon

1CTIVI INGRI DIENTS	
TRIFTHANOLAMINE 2, 4, 5-TRICHLO	-
ROPHENONYACEI ATI	
(Equivalent to 41 of 2-4, 5-1 Acid)	
INTRT INGREDITATE	35.11
TOTAL .	, , , , , , , , , , , , , , , , , , ,
i	i

For control of many kinds of brush and woody plants such as Alder, Aspen. Bucle Chokeberry, Elder, Elderberry, Elm. Hickory, Locust, Poison Ivv. Poison Oak Sumac, Wild Cherry, Wild Grape and Willow, on range lands, rights-of-way, stump land and waste land areas where it is desired to kill or repress growth of woody plants.

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

#### (SLE BACK OR SUDE PANIA FOR OTHER PRECAUTIONS)

On not take internally. May cause skin aritation. Avoid contact with eyes of and alothing. Avoid breathing spray mist

A second place de totale a capt being a television of a proposition of the Second of a captain of the total second of a paper of the definition of the Second of the captain of the definition of the asset of the captain of the television of the captain of the ca

(a) A control of the control of the process of the control of t



BAIRD & McGUIRE, INC.

HOLBROOK, MASSACHUSETTS

A Company of the State of the S

COMMENTS

Complete the trade of the extremal figure