

ACCEPTED  
 June - 27  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 50000

Net Contents

# 4T-Ester Brush Killer

Contains 4 lbs. 2, 4, 5-T Acid Equivalent Per Gallon

ACTIVE INGREDIENTS:	
2-ETHYLHEXYL ESTER OF 2, 4, 5 - TRICHLOROPHENOXYACETIC ACID .....	64.3%
(Equivalent to 44.5% of 2,4,5 - T Acid)	
INERT INGREDIENTS .....	35.7%
TOTAL .....	100.0%

## DIRECTIONS FOR USE

**General:** This product is intended for use in control of brush and woody plants, and should not be used for weed control in growing crops.

**Foliage Spray:** For woody growth less than 6 feet tall, dilute at rate of 3 quarts per 100 gallons of water and spray bark and foliage thoroughly. Spray when plant is in active growing condition. If new growth appears, repeat application.

**Stump Treatment:** For plants 6 feet or higher, cut near ground. Dilute at rate of 3 to 4 quarts per 25 gallons of fuel oil or kerosene and spray bark and cut surfaces.

**Dormant Treatment:** Dilute at the same treatment and spray brush stems and trunks thoroughly to height of 12 to 18 inches.

**Mesquite:** Dilute 1 gallon in 5 gallons of diesel fuel and mix with 13 gallons of water. Apply mixture at rate of 4 gallons per acre when plants have reached the foliage stage of growth. Re-spray 40-60 days after first application.

For control of many kinds of brush and woody plants on range lands, rights-of-way, stump land and waste land areas where it is desirable to kill or repress growth of woody plants, Alder, Blackberry, Cottonwood, Elm, Hazel, Poison Ivy, Raspberry, Sweet Fern, Willow.

## CAUTION

Keep out of reach of children

(See back or side panels for other precautions.)

Do not take internally. May cause skin irritation. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist. Harmful if swallowed.

Avoid spray drift to susceptible plants as this product is toxic to cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Because of difficulty of cleaning sprayers and other equipment used, it is recommended that such equipment not be used for handling and applying other agricultural chemicals. Do not store near fertilizers, acids, insecticides or fungicides.

Vapors from this product may harm plants and animals in proximity.

Prose equipment, air catchers and dip-seal valves should be used when refilling tanks away from water supplies. Containers should be secured by punching holes in them and burying with waste.



U.S. PATENT OFFICE  
 No. 2,541,197

Manufactured by  
 FAIRBANKS MACHINERY, INC.  
 FAIRBANKS, MASS.

## OBSERVE THESE PRECAUTIONS

1. Do not apply directly to, or otherwise permit contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other desirable plants.
  2. Avoid spray drift as even minute quantities of the spray may cause severe injury during both growing and dormant periods. (Coarse sprays are less likely to drift.)
  3. Do not use on lawns or pastures.
  4. Do not contaminate irrigation water or water used for domestic purposes. Excessive amounts of 4T-Ester Brush killer in the soil may temporarily inhibit seed germination and plant growth.
  5. Avoid storing near fertilizers, seeds, insecticides or fungicides.
  6. Do not reuse this container for any other material.
  7. Do not reuse spray rig which has been used for spraying or storing of 2, 4, 5-T solutions.
  8. Use wisely and according to directions. Consult your local agricultural experiment station for specific applications.
  9. Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.
  10. Do not use around the home, recreation area, or similar area.
- This product is toxic to fish. Keep out of streams, rivers or ponds. Apply this product only as specified on this label.