

### WARNING

Avoid inhaling spray.

Coarse sprays are less likely to drift.

Applications by air, ground or hand equipment should be made only when there is no hazard from drift. Spray drift may cause injury to susceptible plants such as vegetables, flowers, grapes or other plants sensitive to minute quantities of herbicide. To avoid injury to desirable plants do not apply other agricultural chemicals with the same equipment used for this herbicide. Do not apply by air in the vicinity of sugar beets.

Do not store near fertilizer, seeds, insecticides, fungicides, feed or foodstuffs.

Excess amounts of this product in soil may temporarily inhibit seed germination or plant growth.

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of waste and containers. Destroy empty container by perforating and bury in a safe place.

Toxic to fish.

Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions. Be sure that the use of this product conforms to all applicable conditions.

### WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability of fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

21517

USDA Registration Number 554-124

NET CONTENTS

GALLONS



# AGSCO 600 HERBICIDE

#### ACTIVE INGREDIENT:

Diethanolamine Salt of		
*2,4-Dichlorophenoxyacetic acid	- - - - -	60.20%
INERT INGREDIENTS	- - - - -	39.80%
		100.00%

\*Acid equivalent 41.00%

Contains 4 Pounds Acid Per Gallon

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

(Additional precautionary statements on side panel)

Harmful if swallowed. Do not get in eyes. Avoid contact with skin or clothing. In case of contact with weed killer, flush with plenty of water for fifteen (15) minutes. For Eyes, get medical attention. Remove and wash contaminated clothing and shoes before re-use.

Protect from temperatures below 0°F.

Manufactured By

AGSCO CHEMICALS, INC. Grand Forks, North Dakota

### DIRECTIONS

This is a selective herbicide used for the control of susceptible weeds in small grains and corn, and for control of certain deep rooted perennials and woody plants in non-crop lands. For best results, apply when soil and climatic conditions are conducive for rapid weed growth.

**WEEDS CONTROLLED:** Mustard, Frenchweed, Arrowhead, Cocklebur, Dandelion, Gumweed, Marsh Elder, Hedge Bindweed, Pigweed, Ragweed, Field Horsetail, Plantain and other broadleaf weeds.

**WHEAT, BARLEY, RYE** — Use 1/2 to 1 pint of this herbicide per acre in sufficient water to obtain good coverage. It is recommended that the lower dosage be used for more susceptible weeds such as common mustard, and the higher dosage level may be necessary for the control of more resistant weeds such as Canada thistle. Make application only between the stooling and joint stages.

**NOTICE:** Do not forage or graze on treated grain fields within two (2) weeks after treatment. Do not feed treated straw to livestock.

**CORN** — Use 1/2 to 1 pint of this herbicide per acre in sufficient water to obtain good coverage. Make application following emergence or majority of weeds but before weeds have damaged the corn. Post emergence application of this herbicide should be made after emergence but before tasseling. Do not apply from tasseling to dough stage. Use drop nozzles when corn is over 8 inches high. Strains and varieties of corn vary in their tolerance to 2,4-D and particular care should be exercised in using either amine or ester formulations. Do not apply to light sandy soil.

**DEEP ROOTED PERENNIALS AND WOODY PLANTS** — In areas such as ditch banks, highway right-of-ways and fence rows, use 1/2 to 3 pints of this herbicide per acre in sufficient water to insure good coverage. In the control of plants such as willows and poison oak, it is necessary to have thorough coverage. Certain weeds and woody plants will form new growth after treatment and repeat applications may be necessary. For best results apply in spring after new leaves appear and plants are growing rapidly.

ACCE  
3-8-71  
554-124

## DIRECTIONS

**AGSCO 666** is a selective herbicide used for the control of susceptible weeds in small grains and corn, and for control of certain deep rooted perennials and woody plants in non-crop lands. For best results, apply when soil and climatic conditions are conducive for rapid weed growth.

**WEEDS CONTROLLED:** Mustard, Frenchweed, Arrowhead, Dandelion, Cocklebur, Gumweed, Marsh Elder, Hedge Bindweed, Pigweed, Ragweed, Wild Buckwheat, Plantain, and other broad-leaf weeds.

**WHEAT, BARLEY, RYE** - Use  $\frac{1}{3}$  to  $\frac{1}{2}$  pint of this herbicide per acre in sufficient water to obtain good coverage. It is recommended that the lower dosage be used for more susceptible weeds such as common mustard, and the higher dosage level may be necessary for the control of more resistant weeds such as Canada thistle. Make application only between the stooling and joint stages.

**NOTICE** - Do not forage or graze on treated grain fields within two (2) weeks after treatment. Do not feed treated straw to livestock.

**CORN** - Use  $\frac{1}{3}$  pint of this herbicide per acre in sufficient water to obtain good coverage. Make application following emergence of majority of weeds but before weeds have damaged the corn. Post emergence application of this herbicide should be made after emergence but before tasseling. Do not apply from tasseling to dough stage. Use drop nozzles when corn is over 8 inches high. Strains and varieties of corn vary in their tolerance to 2,4-D and particular care should be exercised in using ester formulations.

**DEEP ROOTED PERENNIALS AND WOODY PLANTS** - In areas such as drainage ditch banks, highway right of ways and fence rows, use 1 to 2 pints of this herbicide per acre in sufficient water to insure good coverage. In the control of plants such as willows and poison oak, it is necessary to have thorough coverage. Certain weeds and woody plants will form new growth after treatment and repeat applications may be necessary.

EPA Est. No. 554-ND-1

EPA Reg. No. 554-130

NET CONTENTS  GALLONS



# AGSCO 666

## HERBICIDE

A low volatile formulation

ACTIVE INGREDIENT:	2,4-Dichlorophenoxyacetic acid, Isocetyl Ester	94.40%
INERT INGREDIENTS:		5.60%
		100.00%

2,4-Dichlorophenoxyacetic acid equivalent 62.6  
Contains 6 Pounds Acid Equivalent Per Gallon

ACCEPTED

August 16, 1974

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
PD UNDER NO. 554-130

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**

(Additional precautionary statements on side panel)

Harmful if swallowed. Do not get in eyes. Avoid contact with skin or clothing. In case of contact with weed killer, flush with plenty of water for fifteen (15) minutes. For eyes, get medical attention. Remove and wash contaminated clothing and shoes before re-use.

Protect from temperatures below 0°F.

Manufactured By

**AGSCO CHEMICALS, INC.** Grand Forks, North Dakota

## WARNING

Avoid inhaling spray.

Coarse sprays are less likely to drift.

Spray drift or vapors may cause injury to susceptible plants such as vegetables, flowers, grapes or other plants sensitive to minute quantities of herbicide. To avoid injury to desirable plants, do not apply other agricultural chemicals with the same equipment used for this herbicide. Do not apply by air in the vicinity of sugar beets.

Do not store near fertilizers, seeds, insecticides, fungicides or foodstuffs.

Excess amounts of this product in soil may temporarily inhibit seed germination or plant growth.

Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Applications by air, ground or hand equipment should be made only when there is no hazard from drift. Do not contaminate water by cleaning of equipment or disposal of wastes. Be sure that the use of this product conforms to all applicable conditions.

Destroy empty container by perforating and bury in a safe place. Do not burn.

Do not contaminate water used for irrigation or domestic purposes.

## WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**WARNING**

Avoid inhaling spray.

Coarse sprays are less likely to drift.

Applications by air-ground or hand equipment should be made only when there is no hazard to nearby plants. Spray drift may cause injury to susceptible plants such as vegetable, ornamental, flower, shrub, or other plants sensitive to minute quantities of herbicide. To avoid injury to desirable plants, do not apply other agricultural chemicals with the same equipment used for this herbicide. Do not apply spray in the vicinity of water bodies.

Do not use near fertilizers, acids, or other agricultural chemicals, feed or food stuffs.

Excess amount of the product in soil will temporarily inhibit seed germination in plants grown.

Do not contaminate any body of water by direct application, cleaning of equipment, or spillage. Wash and clean containers. Do not empty container by pouring and bury in a safe place.

Toxic to fish.

Excess use may affect the use of fertilizer. State agricultural authorities in each state issue recommendations for application rates. Be sure that the use of the product conforms to all applicable regulations.

**WARRANTY**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. But neither this warranty nor any other warranty of merchantability of fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under abnormal conditions or other conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

21517

USDA Registration Number 554-124

NET CONTENTS

GALLONS



**AGSCO 600  
HERBICIDE**

**ACTIVE INGREDIENT**

Diethanamine Salt of 2,4-Dichlorophenoxyacetic acid 60.20%

**INERT INGREDIENTS** 39.80%

100.00%

Acid equivalent 41.60%

Contains 4 Pounds Acid Per Gallon

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

(Additional precautionary statements on side panel)

Harmful if swallowed. Do not get in eyes. Avoid contact with skin or clothing. In case of contact with weed killer, flush with plenty of water for fifteen (15) minutes. For Eyes, get medical attention. Remove and wash contaminated clothing and shoes before re-use.

Protect from temperatures below 0°F

Manufactured By

**AGSCO CHEMICALS, INC.** Grand Forks, North Dakota

Tr  
bl  
de  
Fe  
co  
  
W  
Cl  
wa  
E  
  
W  
Co  
is  
No  
ch  
wa  
the  
  
NC  
in  
to  
  
CO  
sul  
for  
has  
her  
ins  
not  
eto  
can  
lath  
  
DEI  
are  
row  
cer  
sur  
the  
new  
rec  
app

3-8-71  
554-124

NET CONTENTS

GALLONS



# AGSCO 600 HERBICIDE

## ACTIVE INGREDIENT

Diethanamine Salt of  
2,4-Dichlorophenoxyacetic acid 60.20 %

INERT INGREDIENTS 39.80 %

100.00 %

Acre Equivalent: 41.67

Contains 4 Pounds Acid Per Gallon

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

(Additional precautionary statements on side panel)

Harmful if swallowed. Do not get in eyes. Avoid contact with skin or clothing. In case of contact with weed killer, flush with plenty of water for fifteen (15) minutes. For Eyes, get medical attention. Remove and wash contaminated clothing and shoes before re-use.

Protect from temperature below 0°F

Manufactured By

AGSCO CHEMICALS, INC. Grand Forks, North Dakota

## DIRECTIONS

This is a selective herbicide used for the control of susceptible weeds in small grains and corn, and for control of certain deep rooted perennials and woody plants in non-crop lands. For best results apply when soil and climatic conditions are conducive for rapid weed growth.

**WEEDS CONTROLLED:** Mustard, Fox weed, Am. weed, Cocklebur, Dandelion, Gumweed, Marsh Elder, Hedge Bind weed, Pigweed, Ragweed, Field Horsetail, Plantain and other broadleaf weeds.

**WHEAT, BARLEY, RYE** — Use 1 to 3 pints of this Herbicide per acre in sufficient water to obtain good coverage. It is recommended that the lower dosage be used for more susceptible weeds such as common mustard, and the higher dosage level may be necessary for the control of more resistant weeds such as Canada thistle. Make application only between the following stages:

**NOTICE:** Do not graze or plow in treated grain field within two (2) weeks after treatment. Do not feed treated straw to livestock.

**CORN** — Use 1 to 3 pints of this herbicide per acre in sufficient water to obtain good coverage. Make application following emergence of majority of weeds but before weeds have damaged the corn. Post-emergence application of this herbicide should be made after corn is 6 to 8 inches high but before tasseling. Do not apply from tasseling through stage. Use drop nozzles when corn is over 8 inches high. Strains and varieties of corn vary in their tolerance to 2,4-D and particular care should be exercised in using other amine ester formulations. Do not apply to light soils.

**DEEP ROOTED PERENNIALS AND WOODY PLANTS** — In areas such as ditch banks, highway right of ways and fence rows use 1 to 3 pints of this herbicide per acre in sufficient water to obtain good coverage. In the control of plants such as willows and goose oak it is necessary to have thorough coverage. Certain weeds and woody plants will form new growth after treatment and repeat applications may be necessary. For best results apply in spring after new leaves appear and plants are growing rapidly.

## WARNING

Do not drift

Use hand equipment should be no hazard to nearby plants. Spray susceptible plants with a wetting agent. Do not spray other plants sensitive to this herbicide. To avoid injury to desirable plants, do not spray other agricultural chemicals, or use of this herbicide. Do not spray sugar beets.

Do not use in orchards, vineyards, or on other plants.

Do not use on plants that are sensitive to this herbicide.

Do not use on plants that are sensitive to this herbicide.

Do not use on plants that are sensitive to this herbicide.

## WARRANTY

This product conforms to its chemically stated composition and is reasonably fit for the purpose for which it is intended. It is used in accordance with the directions of use, but neither this warranty of merchantability of purpose, express or implied, extends to any product contrary to label instructions, or under conditions not intended by the seller and buyer.

Registration Number 554-124

3-8-71  
554-124