

GRO-TONE®

Lawn Fungicide Spray

contains Terraclor E.C.+

CONTROLS Brown Patch

APPLY to St. Augustine, Bermuda and Similar Warm Season Grasses

ACTIVE INGREDIENT

Pentachloronitrobenzene (PCNB) . . . . .	24.0%
INERT INGREDIENTS . . . . .	76.0%
TOTAL . . . . .	100.0%

+Terraclor fungicide, Terraclor, registered trademark, Olin Mathieson Chemical Company.

KEEP OUT OF REACH OF CHILDREN

CAUTION


See back panel for additional precautionary information and specific directions for use. Read the label before using.

Manufactured for VIGORO INDUSTRIES, INC., Fairview Heights, IL 62208. GRO-TONE® is a registered trademark of Vigoro Industries, Inc.

NET CONTENTS 1 PINT

EPA Reg. No. 557-1936

EPA Est. No. 6720-FL-1



PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May cause irritation of eyes, skin, nose and throat. Avoid contact with eyes, skin and clothing. Avoid contact with skin by wearing the following protective clothing; gloves, long-sleeved shirt, long pants, socks and shoes. Wash hands thoroughly after using. Avoid breathing vapor or spray mist. Keep container closed. Use with adequate ventilation. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Do not induce vomiting. Get medical attention immediately.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash with soap and warm water.

If in eyes: Rinse thoroughly with clean water. Get medical attention immediately.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water or wetlands by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not graze treated areas. Do not feed clippings to livestock.

PHYSICAL OR CHEMICAL HAZARD

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**Storage:** Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizers seeds, food or feed.

**Product Disposal:** Securely wrap original container in several layers of newspaper and discard in trash. Soak up spilled product in absorbent material before disposal.

**Container Disposal:** Do not reuse empty container. Wrap container and put in trash.

REAR PANEL (Cont'd)

For control of Brown Patch (*Rhizoctonia Solani*) attacking St. Augustine, Bermuda and similar warm season grasses, apply 2 tablespoons per gallon of water to 30 square feet (5 feet x 6 feet) or 1 pint in 15 to 20 gallons water to cover 500 square feet (20 feet x 25 feet). For easy application, use hose end type sprayer designed to deliver 15 to 20 gallons dilute spray while emptying quart jar. Pour 1 pint concentrated liquid into jar and fill to top with water and apply to 500 square feet. Apply in spring or fall at first sign of disease and repeat in 3 to 4 weeks if necessary. Sprinkle after application to wash spray into soil surface. Follow recommended rate. Excess may cause a temporary yellowing.

CONDITIONS OF SALE: All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.