

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR COMPLETE CAUTION STATEMENT.



ROCKLAND

RESIDUAL FLY SPRAY



ACCEPTED
11-29-67
572-124
FOR THE FEDERAL REGISTER
FOR THE FEDERAL REGISTER
ED UNDER THE FEDERAL REGISTER
TO ATTACHED COMMENTS.

KILLS

see add'l
Flies and Livestock Pests
On ~~Animals~~, In Farm Buildings
4-6 week Residual Fly Control *see add'l*

| | | |
|---|-------|--------|
| ACTIVE INGREDIENTS: | | |
| *Ronnel (0,0-dimethyl 0-2,4,5-trichlorophenyl phosphorothioate) | 24.0% | |
| Aromatic Petroleum derivative solvent | 68.0% | |
| INERT INGREDIENTS: | 8.0% | |
| | | 100.0% |

*From technical grade; not suitable for internal use in animals.

U. S. Pat. No. 2599516

Rockland Chemical Co., Inc.
Passaic Avenue, West Caldwell, N. J.

DIRECTIONS FOR USE

ROCKLAND RESIDUAL FLY SPRAY mixes readily with water or oil to be used as directed as residual fly spray or as a livestock insect spray.

RESIDUAL FLY SPRAY — Farm Buildings, including Poultry Houses, Livestock Shelters, Milk Rooms, Dairy and Hog Barns. For 4-6 weeks control, mix 1 qt. ROCKLAND RESIDUAL FLY SPRAY with 6 gallons of water (5 oz. to 1 gallon water). Add 3 ounces of sugar for added effectiveness.

Spray walls, ceilings, partitions to obtain complete coverage with no run-off. Use 1 gallon of diluted spray per 500-1000 sq. ft. To prevent contamination, keep milk and milking equipment covered while spraying.

For Caged Layers on Wire, mix 1 qt. ROCKLAND RESIDUAL FLY SPRAY with 3 gallons of water. Spray poultry droppings beneath cages.

LIVESTOCK INSECT SPRAY — Dilute with water only. Spray entire animal to wet coat thoroughly. Repeat as necessary but not oftener than every 2 weeks.

Cattle, Calves; to control Hornflies, ~~Screech-worms~~, use 1 qt. with 12 1/2 gallons water. To control Lice, use 1 qt. with 25 gallons water. To control Lone Star Ticks and Winter Ticks, use 1 qt. with 8 gallons water.

Sheep, Goats, Lambs and Kids; to control ~~Screech-worms~~, Wool Maggots, Hornflies, use 1 qt. with 12 1/2 gallons water. To control Lice and Sheep Keds, use 1 qt. with 25 gallons water.

Hogs; to control Lice, use 1 qt. with 25 gallons water.

Cattle Backrubbers — for Hornflies and to aid in control of Face Flies if backrubbers are placed so animal can rub face. Mix 1 qt. ROCKLAND RESIDUAL FLY SPRAY with 7 gallons of diesel, or similar oil. Saturate fibre; recharge as necessary. Do not mix with used motor oil.

WARNING: Do not treat hogs within 6 weeks, cattle within 8 weeks, sheep and goats within 12 weeks of slaughter. Do not treat lactating dairy animals, or dry dairy animals within 21 days of the lactating period. Do not allow spray to come in contact with food, feed or water. Remove lactating dairy animals from buildings during application of residual fly spray. *see add'l*

NOTE TO PHYSICIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

CAUTION: INJURIOUS TO THE EYES

IRRITATING TO THE SKIN

HARMFUL IF SWALLOWED

Do not get in the eyes. Avoid prolonged or repeated contact with the skin.

Avoid prolonged breathing of spray mists

Avoid contamination of food, feedstuffs, and water supplies.

Keep material out of reach of children and animals.

Keep away from heat and open flame.

Keep container closed. Wash thoroughly with soap and water after handling.

COMBUSTIBLE LIQUID.

WARRANTY: Seller makes ~~no~~ warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not. *Replace*

When used as directed on label and in accordance with good practices, this material complies with the Food and Drug Law and other Federal Laws. *Remove*

ONE QUART

US Pat Reg No 572-124