## Renomyl Turf Fungicide Granules



to base of plants.

For Large Brown Patch, spread contents of bag over 3,000 sq. ft. of turf. For Dollar Spot, spread contents of bag over 6,000 sq. ft. of turf.

Apply at earliest appearance of disease and continue at 10 to 14 day intervals, as long as needed, or 5 to 7 days when conditions are unusually favorable for development of disease. Do not graze or feed clippings from treated areas to livestock.

For Stripe Smut, spread contents of bag over 1,000 sq. ft. Make single application in early Spring or in October.

## SUGGESTED SPREADER SETTINGS

	•	1,000 Sq. Ft. Coverage		3,000 Sq. Ft. Coverage	
CYCLONE	4¼4½	6-6¼	GREENVIEW	7¾8¼ .	10-101/2
SCOTT DROP	66½	11½-12	SCOTT ROTARY	DE	N1/2-01/2

These spreader settings are meant as a guide. They are in no sense a guarantee of exact application, inasmuch as age, wear of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

## STORAGE AND DISPOSAL

STORAGE: Never allow this product to become wet during storage. Store unused material in original container in a locked storage area out of reach of children and animals. Keep container tightty closed. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL: Completely empty beg into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-256 #B25612 E.P.A. Est. 9198-OH-1

NET WEIGHT 12 LBS.
Treats 3,000 Sq. Ft. for Large Brown Patch.

Treats 1,000 Sq. Ft. for Stripe Smut.

Treats 6,000 Sq. Ft. for Dollar Spot.

## MANUFACTURED FOR