

UNDER THE FEDERAL

# 3% DODINE DUST

FUNGICIDE DUST - CONTAINS 3% CYPREX<sup>®</sup>

**ACTIVE INGREDIENTS:**

Dodine (n-dodecylguanidine acetate) .....	3.0%
INERT INGREDIENTS: .....	97.0%
	100.0%

A FUNGICIDE DUST FORMULATION SUGGESTED FOR USE BY GROUND DUSTER OR AIRPLANE, FOR CONTROL OF APPLE SCAB, PEAR SCAB AND SOUR CHERRY LEAF SPOT.

## CAUTION: Keep Out of Reach of Children

### MATERIAL CAUSES EYE IRRITATION

In case of contact with eyes, flush with plenty of water for at least fifteen minutes; obtain medical attention.

**DO NOT APPLY DODINE IN COMBINATION WITH LIME OR CHLOROBENZILATE.**

Avoid contamination of feed and foodstuffs.

Do not apply when conditions favor drift from area treated.

Care should be taken to avoid application on peaches, plums and bearing sweet cherries since severe foliage or fruit injury may result. Do not graze cover crops in treated orchards. Consult your State Agricultural Experimental Station or Extension Service for proper timing of applications.

## NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

## RECOMMENDATIONS FOR USE

**APPLES:** Apple Scab — Apply 25 to 40 lbs. of 3% Dodine Dust per acre at 5 to 7 day intervals or as needed to maintain scab control from pre-bloom through first cover applications. Use 40 lbs. of Haviland's 2% Dodine Dust per acre in later cover applications. Do not apply within 5 days of harvest. Dodine treated apples should not be used in the manufacture of apple pomace for use in livestock feeds. Such use may result in illegal residues in milk or meat.

**PEARS:** Pear Scab — Western States Only — Apply 25 to 40 lbs. of 3% Dodine Dust per acre at 5 to 7 day intervals or as needed to maintain scab control from pre-bloom through first cover applications. Continue applications at 10 to 14 day intervals or as needed. Use 40 lbs. of Haviland's 2% Dodine Dust per acre in later cover applications. Do not apply within 5 days of harvest.


**SOUR CHERRIES:** Leaf Spot — Apply 50 to 60 lbs. per acre of 3% Dodine Dust at petal-fall or when first leaves unfold. Repeat application thereafter at 7 to 10 day intervals. Use the shorter interval during periods of rapid growth and/or frequent rains. If conditions favor leaf spot, continue applications through harvest. Post-Harvest Sprays — Apply 50 lbs. per acre of 3% Dodine Dust immediately after harvest and as needed.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

\*CYPREX<sup>®</sup> is the registered Trademark of the American Cyanamid Company.

USDA Reg. No. 595 265

Net Wt. 50 Lbs.



# HAYILAND

1345 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502