

THIODAN MANEB DUST

(INSECTICIDE-FUNGICIDE DUST 3% - 7%)

ACTIVE INGREDIENTS:

Endosulfan (Hexachlorohexahydromethano-2, 4, 3-benzodioxathiepin oxide)	3.0%
Maneb (Manganese ethylenebisdithiocarbamate).	7.0%
INERT INGREDIENTS	90.0%
	100.0%

Thiodan® is a registered trademark of Farbwerke Hoechst A.G. for the active ingredient commonly known as endosulfan.

DIRECTIONS

The following directions are furnished by us and are believed to be reliable but in no way guaranteed. Make first application when insects appear or disease threatens and repeat weekly if necessary. Consult your State Agricultural Experiment Station or Extension Service for specific timing and dosage rates in your area.

BEANS: To control Mexican Bean Beetles, Downy Mildew and Rust, apply 25 to 30 lbs. per acre. Do not apply after pods begin to form. Do not feed treated threshings to livestock.

BROCCOLI, CABBAGE AND CAULIFLOWER: To control Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth Larvae, Alternaria Leaf Spot and Downy Mildew. Do not apply on cauliflower after edible parts begin to form and do not apply to broccoli or cabbage within 7 days of harvest. Apply 20 to 25 lbs. per acre.

CUMCUMBERS, MELONS AND SQUASH: To control Aphids, Anthracnose and Downy Mildew apply 25 to 30 lbs. per acre. Apply as needed up to harvest.

HEAD LETTUCE: To control Aphids, Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth Larvae and Downy Mildew apply 25 to 33 lbs. per acre. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove outside leaves at harvest. Do not feed crop refuse to livestock.

PEPPERS AND TOMATOES: To control Aphids, Colorado Potato Beetle, Flea Beetles, Anthracnose and Early and Late Blight apply 15 to 20 lbs. per acre. Do not apply within 4 days of harvest. When applying in

greenhouse, wear a mask or respirator passed by the U.S. Bureau of Mines for Thiodan protection. To avoid damage, do not use on tender, young greenhouse plants.

POTATOES: To control Aphid, Colorado Potato Beetle, Flea Beetles, Leafhoppers and Early and Late Blight apply 30 to 35 lbs. per acre. Apply as needed up to harvest. Do not plant root crops other than carrots, potatoes or sweet potatoes as followup crops.

CAUTION: Keep Out of Reach of Children

HAZARDOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN!

Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water, for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. During commercial or prolonged exposure in spray mixing and loading operations, wear a mask or respirator of a type passed by the U.S. Bureau of Mines for endosulfan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed or foodstuffs.

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning of equipment or disposal of wastes. Keep out of lakes, streams or ponds.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. Reg. No. 595-275

ACCEPTED
6-13-72 with comments
595-275
Net Wt. 50 Lbs.



HAYLAND

1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502

ACCEPTED WITH COMMENTS