

captan diazinon dust no. 752

ACTIVE INGREDIENTS:

Captan [N- [(trichloromethyl) thio] -4-cyclohexene -1, 2-dicarboximide]	7.5%
[O, O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) thiophosphate]	2.0%
INERT INGREDIENTS:	90.5%
	100.0%

CAUTION:

Keep Out of Reach of Children

Harmful if swallowed or inhaled! If accidentally swallowed, empty the stomach by using an emetic or by mechanical means. Avoid breathing dust. Wear a respirator or mask of a type approved by the U.S. Bureau of Mines. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before eating or smoking. Do not contaminate feed or foodstuffs.

Do not contaminate water by cleaning of equipment or disposal of waste.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. This product is toxic to fish, birds, and other wildlife. Birds and wildlife in treated areas may be killed.

Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS

GENERAL APPLICATION: Apply dust weekly at the rate listed. Use the higher dosages at shorter intervals, under severe infection or pest conditions.

APPLES: To control apple scab, Brooks fruit spot, sooty blotch, fly speck, black rot, bitter rot, black pox, and botryosphaeria, apply at the rate of 35 to 50 lbs. per acre. To control aphids, mites, San Jose scale crawlers, and Forbes scale crawlers, apply at the rate of 45 to 75 lbs. per acre. Begin application at first cover, repeating as needed to maintain adequate coverage. Do not apply within 14 days of harvest. Slight russeting may occur on some varieties of apples.

CHERRIES: To control brown rot, leaf spot, black cherry aphids, and cherry fruit flies, apply at the rate of 25 to 50 lbs. per acre. To control San Jose scale crawlers, apply at the rate of 45 to 75 lbs. per acre. Begin application at petal fall and repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest.


PEACHES: To control brown rot, peach scab, and aphids, apply at the rate of 30 to 50 lbs. per acre. To control San Jose scale crawlers, apply at the rate of 45 to 75 lbs. per acre. Begin application at petal fall and repeat at 7 to 10 day intervals or as necessary to maintain adequate control. Do not apply within 20 days of harvest.

STRAWBERRIES: To control botrytis fruit rot, aphids, and mites, apply at the rate of 30 to 40 lbs. per acre. Begin application when pests or disease are first noted, and repeat at 7 day intervals or as needed. Do not apply within 5 days of harvest. For insect control, dust should be directed into the crowns of the plant and plants sufficiently agitated to insure thorough coverage.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. Reg. No. 595-278



Net Wt. 50 Lbs.



HAVILAND

1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502

ACCEPTED WITH COMMENTS