

ACCEPTED

JUN 14 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 35217 SUBJECT
TO ATTACHED COMMENTS.

6% Dodine Dust
(fungicide dust - contains 6% Cyprex*)

ACTIVE INGREDIENTS

Dodine (n-dodecylguanidine acetate) 6.0%

INERT INGREDIENTS 94.0%

Total..... 100.0%

A FUNGICIDE DUST FORMULATION SUGGESTED FOR USE BY GROUND DUSTER OR AIRPLANE, FOR CONTROL OF APPLE SCAB, PEAR SCAB AND SOUR CHERRY LEAF SPOT.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
CONCENTRATED MATERIAL CAUSES EYE IRRITATION**

In case of contact with eyes, flush eyes with plenty of water for at least fifteen minutes; obtain medical attention.

DO NOT APPLY DODINE IN COMBINATION WITH LIME OR CHLOROBENZILATE.

Avoid contamination of feed and foodstuffs.

Avoid drift of dust on adjacent crops.

Care should be taken to avoid application on cherries, plums and bearing sweet cherries since severe foliage or fruit injury may result.

Do not graze cover crops in treated orchards.

Consult your State Agricultural Experiment Station or Extension Service for proper timing of applications.

NON-WARRANTY

The use or handling of this material being beyond our knowledge and control we do not make any warranty expressed or implied as to the effects of such use whether or not in accordance with directions or claimed to be so.

RECOMMENDATIONS FOR USE

APPLES: Apple-Scab - Apply 15 to 20 lbs. of 6% Dodine dust per acre at 5 to 7 day intervals or as needed to maintain scab control from pre-bloom through first cover applications. Use 40 lbs. of "Haviland's 2% Dodine Dust" per acre in later cover applications. Do not apply within 5 days of harvest. Dodine treated apples should not be used in the manufacture of apple pomace for use in livestock feeds. Such use may result in illegal residues in milk or meat.

PEARS: Pear Scab - Western States Only -- Apply 15 to 20 lbs. of 6% Dodine Dust per acre at 5 to 7 day intervals or as needed to maintain scab control from pre-bloom through first cover applications. Continue applications at 10 to 14 day intervals or as needed. Use 40 lbs. of "Haviland's 2% Dodine Dust" per acre in later cover applications. Do not apply within 5 days of harvest.

HAVILAND AGRICULTURAL CHEMICAL COMPANY
GRAND RAPIDS, MICHIGAN



^{Sour}
SOUR CHERRIES: Leaf Spot - Apply 25 to 30 lbs. per acre of 6% Dodine Dust at petal-fall or when first leaves unfold. Repeat application thereafter at 7 to 10 day intervals. Use the shorter interval during periods of rapid growth and or frequent rains. If conditions favor leaf spot, continue applications through harvest. Post-Harvest Sprays - Apply 25 lbs. per acre of 6% Dodine Dust Immediately after harvest and as needed.

CYPREX* is the registered Trademark of the American Cyanamid Company.

USDA Reg. No. _____ Net Wt. 50 Lbs.

HAVILAND AGRICULTURAL CHEMICAL COMPANY
GRAND RAPIDS, MICHIGAN