

3% METHOXYCHLOR COPPER DUST

(Insecticide-Fungicide)

ACTIVE INGREDIENTS:

*Methoxychlor, Technical	3.0%
Copper, metallic (from 2-5 micron Tribasic Copper Sulfate) ..	5.0%
INERT INGREDIENTS:	<u>92.0%</u>
Total	100.0%

*Equivalent to 2.64% 2,2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 0.36% other isomers and related compounds.

CAUTION: Keep Out Of Reach Of Children

Do not exceed amount given nor apply closer to harvest than is recommended. Wash treated produce thoroughly before eating, marketing or processing. Avoid inhalation and contamination of foodstuffs.

Toxic to fish. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-317

9-7-71
595-317

Net Wt. 50 Lbs.

DIRECTIONS

The following directions are furnished by us and are believed to be reliable. Make thorough, full-coverage applications as insects or disease threatens. Repeat applications at 5 to 7 days as necessary. Rates suggested below are in lbs. per acre. Do not use excessive amounts or closer to harvest than recommended below. Consult State Agricultural Experiment Station for proper timing of applications.

<u>CROP</u>	<u>INSECT</u>	<u>DISEASE</u>	<u>RATE</u>	<u>LIMITATIONS</u>
<u>BEETS</u>	Flea Beetles Leafhoppers	Early & Late Blight Downy Mildew Leaf Spot	60-70	7 days*
		*14 days if tops are used as feed or food.		
<u>CABBAGE, CAULIFLOWER, & BROCCOLI</u>	Imported Cabbageworm Flea Beetles	Downy Mildew Leaf Spot	40-45	Cabbage 3 days Cauliflower 7 days Broccoli 14 days
<u>CARROTS</u>	Leafhoppers	Downy Mildew Leaf Spot	30-40	14 days
<u>CUCUMBERS, MELONS, & SQUASH</u>	Cucumber Beetles Leafhoppers Melon Worm Pickleworm	Angular Leaf Spot Anthracnose Bacterial Wilt Downy Mildew Powdery Mildew	30-40	Cucumber 7 days Melons 7 days Squash 1 day
<u>TOMATOES, EGGPLANT, & PEPPERS</u>	Flea Beetle Leafhoppers	Anthracnose Early & Late Blight Septoria Leaf Spot Leaf Mold	60-70	7 days
<u>POTATOES</u>	Leafhoppers	Early & Late Blight	30-45	0

3% METHOXYCHLOR 5% COPPER DUST

(Insecticide-Fungicide)

ACTIVE INGREDIENTS:

*Methoxychlor, Technical 3.0%
Copper, metallic (from 2.5 micron Tribasic Copper Sulfate) .. 5.0%

INERT INGREDIENTS: 92.0%

Total 100.0%

DIRECTIONS

The following directions are furnished by us and are believed to be reliable. Make thorough, full-coverage applications as insects or disease threatens. Repeat applications at 5 to 7 days as necessary. Rates suggested below are in lbs. per acre. Do not use excessive amounts or closer to

Equivalent to 2.64% 2,2-bis (p-methoxyphenyl) 1,1,1-trichloroethane and 0.36% other isomers and related compounds.

Station for proper timing of applications.

CAUTION: Keep Out Of Reach Of Children

Do not exceed amount used per acre, except to harvest fruit as recommended. Wash treated produce thoroughly before eating, marketing or processing. Avoid inhalation and contamination of foodstuffs.

For information on the use of this product, contact your local agricultural agent or the manufacturer. For information on the use of this product, contact your local agricultural agent or the manufacturer.

For information on the use of this product, contact your local agricultural agent or the manufacturer.

NON-WARRANTY

This product is sold as a fungicide and insecticide. It is not a guarantee of performance. The user assumes all responsibility for the proper use of this product. The manufacturer is not liable for any damage or loss resulting from the use of this product.

CROP	INSECT	DISEASE	RATE	LIMITATIONS
BEETS	Beet Beetles Leafhoppers	Early & Late Blight Late Blight Fusarium Wilt Leaf Spot	10-20	Do not use on leafhoppers
CABBAGE CAULIFLOWER & BROCCOLI	Beet Beetles Leafhoppers	Early & Late Blight Late Blight Fusarium Wilt Leaf Spot	10-20	Do not use on leafhoppers
CARROTS	Leafhoppers	Early & Late Blight Late Blight Fusarium Wilt Leaf Spot	10-20	Do not use on leafhoppers
CUCUMBERS MELONS & SQUASH	Beet Beetles Leafhoppers	Early & Late Blight Late Blight Fusarium Wilt Leaf Spot	10-20	Do not use on leafhoppers
POTATOES	Leafhoppers	Early & Late Blight Late Blight Fusarium Wilt Leaf Spot	10-20	Do not use on leafhoppers