

15% ZINEB DUST

ACTIVE INGREDIENTS:

*Zineb (zinc ethylene bisdithiocarbamate)	15%
INERT INGREDIENTS:	85%
Total	100%

*Metallic Zinc equivalent 3.54%

DIRECTIONS

CHERRIES (sour) . . . To control leaf spot, shotholes:

Apply dust at the rate of 12 to 14 lbs. per Acre. Make first application at petal fall and repeat application 10 to 20 days later. A third application should be made immediately after harvest. Do not apply dust within 7 days before harvest.

MUSHROOM . . . To control brown spot or dry bubble (Verticillium), greenmold (Trichoderma), mildew, soft rot, cobweb (Dactylium):

Apply dust at the rate of 1/2 to 1 lb. per 4,000 sq. ft. Begin dusting soon after casing soil is applied and repeat at 3-7 days until small buttons begin to form. After buttons begin to form, apply dust only to casing soil between "breaks". Do not dust buttons.

CAUTION:

Keep Out of Reach of Children

May cause irritation of nose, throat and skin. Do not breathe dust. In case of prolonged or frequent use, wear clean, protective clothing and bathe after each exposure. In case of skin contact, wash with water; for eyes, wash copiously with water and get medical attention. Keep containers closed. Store in a dry place. Keep away from fire and sparks.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-318

NET WT. 50 LBS.



HAYILAND

REGISTERED, N.W. GRAND RAPIDS, MICHIGAN, 1950.