


E-Z-FLO CAPTAN 7.5 DUST

CAUTION: KEEP OUT OF THE REACH OF CHILDREN



ACTIVE INGREDIENT :

Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)

INERT INGREDIENTS:

	7.50%
	92.50%
Total	100.00%

DIRECTIONS FOR USE

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Avoid contact with skin. Avoid inhaling dust. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift of dust from crop area treated.

DIRECTIONS FOR USE

For best results apply when air is calm. A good hand or power dusting machine should be used to secure complete coverage of plants with a light film of dust. Begin application prior to or at first sign of infection. Repeat applications as necessary. Keep new growth protected. Captan should not be used closely following oil sprays or solvent formulations of parathion.

FRUIT CROPS

APPLES- Bitter rot, Black rot, Frog-eye leaf spot, Black pox, Botryosphaeria: Apply 27 to 50 pounds/acre. Apply in late cover and pre-harvest periods. For frog-eye leaf spot also use 27 to 53 pounds/acre in pre-bloom and blossom periods. Brooks fruit spot, Flyspeck: Use 15 to 50 pounds/acre. Apply in cover periods. Scab: For primary infection use 27 to 106 pounds/acre. Apply in pre-bloom, blossom, petal fall, and first cover periods. For secondary infection use 15 to 33 pounds/acre. Apply in cover periods. Sooty blotch: Use 15 to 28 pounds/acre. Apply in cover periods. Note: Red Delicious and other sensitive varieties may be injured by early season applications.

CHERRIES- Botrytis rot: Apply 80 pounds/acre. Apply at petal fall, shuck shed, and in cover sprays. Brown rot: For fruit rot, apply 15 to 66 pounds/acre in pink, full bloom, petal fall, shuck shed, and cover periods. Leaf spot: Apply 40 to 53 pounds/acre. Apply in pink, bloom, petal fall, and first cover periods and in 1 or 2 post-harvest foliage applications.

PEACHES- Brown rot, Rhizopus, Scab: Apply 26 to 66 pounds/acre. Apply in pink, full bloom, shuck shed, cover, and pre-harvest periods.

PEARS- Fruit spot, Scab (primary infection): Apply 27 to 40 pounds/acre. Apply in pink, petal fall, and first two cover periods. Do not use on D'Anjou pears. Russetting may be produced on Bosc pears.

PLUMS, PRUNES- Brown rot: Apply 27 to 66 pounds/acre. Apply at red bud, 75.0 percent bloom, petal fall, and in cover applications as necessary.

STRAWBERRIES- Botrytis rot (gray mold), Leaf spots: Apply 27 to 53 pounds/acre. Begin at new growth. Repeat at 7 day intervals through picking and after harvest.

VEGETABLE CROPS

BEANS- Anthracnose, Downy mildew, Rust: Apply 27 to 40 pounds/acre. Begin before disease appear or at first sign of disease. Repeat at 4 to 10 day intervals or as necessary to maintain control. Damping-off, Root rot: Apply 67 to 80 pounds/acre, worked into top 3 to 4 inches of soil at planting time.

CANTALOUPE, CUCUMBERS, HONEYDEW MELONS, PUMPKINS, SQUASH (summer and winter), WATERMELONS- Angular leaf spot, Anthracnose, Downy mildew: Apply 15 to 50 pounds/acre. Begin at first sign of disease. Repeat at 5 to 7 day intervals.

CARROTS- Alternaria, Cercospora blights, Septoria leaf spot: Apply 30 to 40 pounds/acre. Begin when disease appear. Repeat at 6 to 10 day intervals.

CELERY- Damping-off, Pink rot: Apply 26.6 pounds/acre. Apply 44 to 86 pounds/acre to plant beds when all seeds have germinated. Repeat as necessary.

POTATOES (preharvest)- Early and late blights: Use 24 to 40 pounds/acre. Begin when plants are 2 to 4 inches high or at first sign of disease. Repeat at 5 to 7 day intervals.

TOMATOES (field and greenhouse)- Anthracnose, Early and late blights, Gray leaf spot (Stemphylium), Septoria leaf spot: Apply 27 to 66 pounds/acre. Begin at first sign of disease. Repeat at 5 to 7 day intervals. Damping-off, Root rot: Apply 66 to 100 pounds/acre. Preplanting soil treatment. Work into upper 3 to 4 inches of soil.


ORNAMENTAL CROPS

GRASSES (ornamental in nonpastured areas only)- Leaf spots, Damping-off, Brown patch, Copper spot, Melting out, Seedling blights, and Brown spot on St. Augustine grass: Apply 1.33 to 2.66 pounds/1000 square feet. Begin when growth starts in spring. Repeat at 7 to 14 day intervals throughout season. Do not graze treated areas. Do not feed clippings to livestock.

HOLLYHOCK, LILAC, SNAPDRAGON, SPIREA- Anthracnose: Apply 28 pounds per acre. Apply before disease appears. Repeat at weekly intervals during warm, wet weather.

ROSES- Black spot, Botrytis blossom blight: Apply 30 to 40 pounds/acre. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals; and more frequently during frequent rains and heavy dews.

STOCKS- Botrytis rot: Apply 54 to 66 pounds/acre. Apply when disease appears. Repeat at 7 day intervals or as needed.



EPA Est 635-MI-1;OH-1

EPA Reg. No. 635-194

Net Contents and Manufacturer
On Container