
635-268



## E-Z FLO ROTENONE 1 D

## CAUTION: KEEP OUT OF REACH OF CHILDREN

 ACTIVE INGREDIENTS:
 1.0%

 Rotenone:
 2.0%

 Other Cube Resins:
 97.0%

 INERT INGREDIENTS:
 100.0%

## CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Destroy Container When Empty. Never Re-use.

51-29

EPA Est. 635-MI-1; ON-1

E.P.A. Reg. No. 635-268

Not C

## DIRECTIONS FOR USE

Dust when insects first appear and repeat applications as often as is necessary to control. Always repeat dusting after a rain. Do not apply within 1 according to the rest.

BEANS: Black bean aphid, cucumber beetle, Mexican bean beetle- Use 30th, per acre.

BROCCOLI, CABBAGE: Cabbage looper, imported cabbageworm - Use 15 lbc per acre. Apply when first signs of worm feeding appear and repeat as necessary

BRUSSELS SPROUTS, CAULIFLOWER: Cabbage aphid, cabbage looper, cross-striped cabbageworm, flea beetles, young harlequin bug, imported cabbageworm- Use 15 lbs. per acre.

CARROTS: Green peach aphid- Use 15 lbs. per acre.

CELERY: Green peach aphid- Use 15 lbs. per acre.

COLLARDS: Cabbage looper, young harlequin bug, imported cabbageworm- Use 15 lbs. per acre.

CUCUMBERS, MELONS, SUMMER SQUASH, WINTER SQUASH and PUMPKINS: Aphids— Use 15 lbs. per acre. Apply when aphids first appear and repeat as required. Treat both upper and lower leaf surfaces thoroughly.

PEPPERS: Green peach aphid - Use 15 lbs. per acre. Apply when insects first appearant repeat as required.

EGGPLANTS: Green peach aphid, flea beetle, and Colorado potato beetle-Use 30 lbs. per acre. Dust plants thoroughly, particularly the undersides of the leave

POTATOES: Colorado potato beetle, potato leafhopper, aphids- Use 30 lbs. perac: Apply when insects first appear and repeat as required.

SPINACH: Green peach aphid- Use 15 lbs. per acre.

TOMATOES: Green peach aphid, Colorado potato beetle, flea beetle-Use 30 lbs. per acre. Apply when insects first appear and repeat as required. Dust plants thoroughly, particularly the undersides of the leaves.