

CRABGRASS CONTROL: TIMING IS IMPORTANT. To prevent crabgrass apply Chlordane 45% before seeds germinate. IN MOST AREAS IT MUST BE APPLIED BEFORE APRIL 1ST. It also has long lasting residual action in the soil. A single application will usually provide protection for the entire growing season. **Do not use on newly seeded lawns or turf.**


Chlordane 45%, used as directed, is harmless to established lawns. Chlordane emulsions are readily applied by means of a pressure sprayer, jar and garden hose attachment, or sprinkling can. Apply 1½ quarts in sufficient water (3 to 10 gallons) to drench uniformly 1,000 sq. ft. of lawn or turf. Immediately after application, wash Chlordane off grass onto soil surfaces. Do not allow children or pets on treated area until Chlordane has been watered into the turf and the treated area has dried completely.

WARNING: May be fatal if swallowed! Avoid inhaling spray. Avoid contact with eyes or clothing. In case of contact with skin, wash with soap and water.

CAUTION: Use ONLY at dilutions as directed. Do not contaminate any body of water nor apply to any area not specified on label. Toxic to fish and wildlife. Birds feeding in treated areas may be harmed. Do not use on animals foods, feed crops, or in dairy barns.

Non-Warranty: Buyer assumes all risk in use or handling, whether in accordance with directions or not.

DE 42730



1 PINT

CHLORDANE 45%

KILLS

**DORMANT CRABGRASS, TERMITES,
ROACHES, ANTS, LAWN INSECTS
SPRAY AWAY CRABGRASS & LAWN INSECTS**

ACTIVE INGREDIENTS:

Technical Chlordane*	45%
Petroleum Distillate	49%

INERT INGREDIENT: 6%

*Equivalent to 27.0% octachloro-4,7-methano-tetrahydroindane and 18.0% related compounds.

WARNING: Keep out of reach of children.

See back panel for additional cautions.

LOS ANGELES THOMPSON CHEMICALS CORPORATION ST. LOUIS

DIRECTIONS FOR USE

FOR LAWN INSECTS: White Grubs, Japanese Beetle Larvae, Asiatic Garden Beetle Larvae, Sod Webworms, Lawn Moths, Wireworms, Cutworms, Chiggers, Ticks, Armyworms, and Ants in lawns. As an insecticide, dilute 2 to 4 fl. ozs. (4 to 8 tbs.) with 3 gals. of water to treat 500 sq. ft. of lawn. For sandy soil, use 4 fl. ozs. (½ cup); for solid type soil, use 8 fl. ozs. (1 cup). It is important to water the lawn thoroughly after applying Chlordane. Note **CAUTIONS** concerning children, pets and birds.

TERMITES: Mix 2 quarts of Chlordane 45% into 25 gals. of water. Apply this emulsion at the dosage of 1 gal. per linear foot to a trench 1 to 2 feet deep and 8 to 12 inches wide dug next to the foundation. Near deep foundations, trench should be 30 inches deep, but not below the level of the footing. Apply directly to infested wooden structures. Inspect area yearly to determine possible reinfestation.

ROACHES, SILVERFISH, SPIDERS, ANTS, IN AND AROUND DWELLINGS: Dilute 1½ fl. ozs. (3 Tablespoons) with 1 qt. of water. Thoroughly spray infested cracks and other hiding places and adjacent exposed surfaces where these pests may occur. Spot treat only. Do not contaminate food, drinking water or utensils. Repeat as necessary.

MADE IN U.S.A. USDA Reg No 642-129

4-7-67
642-129