


10 MALATHION DUST

| | | |
|--------------------|--------|------|
| ACTIVE INGREDIENT: | | |
| Malathion * | ----- | 10% |
| INERT INGREDIENTS: | ----- | 90% |
| | TOTAL: | 100% |

* 0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.



SUGGESTIONS FOR USE:

Use 10% MALATHION DUST for the control of boll weevil, desert spider mite, cotton aphid, cotton leafworm and lygus bug on cotton.

The usual rate of application is 10-25 pounds per acre. However, because local conditions and plant size largely determine the rate, frequency and effectiveness of application, it is recommended that Federal or State agricultural authorities be consulted for specific uses.

NET CONTENTS _____ POUNDS

E.P.A. Reg. No. 655-550

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SPECIFIC CAUTION STATEMENTS ON SIDE PANEL.

CAUTION: AVOID PROLONGED BREATHING OF DUST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN; WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. HIGHLY TOXIC TO FISH. DO NOT CONTAMINATE ANY BODY OF WATER NOR APPLY WHEN CONDITIONS FAVOR DRIFT FROM THE CROP AREA TREATED.

BURN OR BURY CONTAINERS; NEVER REUSE. STAY OUT OF SMOKE OR FUMES.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.



Manufactured by

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105
Office: 363 7th Avenue
New York, N.Y. 10001