

FIRST AID: If swallowed, give a glass of warm water and induce vomiting by touching back of throat with a finger. Repeat until vomit fluid is clear. Call a physician immediately. Symptoms of Warfarin poisoning may occur after repeated skin contact. If on skin or in eyes flush immediately with plenty of water. For eyes, see a physician.

NOTE TO PHYSICIAN: When a human has been known to have ingested Warfarin, monitoring of plasma prothrombin activity is indicated. If victim is hemorrhaging, intravenous and oral doses of vitamin K₁ are indicated, along with blood transfusions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Fatal if swallowed or absorbed through skin. Do not get in eyes, on skin, or on clothing. This material can reduce the clotting ability of blood and cause internal bleeding.

ENVIRONMENTAL HAZARDS

Keep away from wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

SODIUM WARFARIN SOLUTION

A RODENTICIDE CONCENTRATE FOR FORMULATING USE

ACTIVE INGREDIENTS:

Sodium Warfarin (Na 3-(alpha acetonyl-benzyl)-4-hydroxycoumarin)	42.85%*
INERT INGREDIENTS	57.15%
TOTAL	100.00%

* Equivalent to 40.0% Warfarin

KEEP OUT OF REACH OF CHILDREN



E.P.A. REG NO. 655-590 E.P.A. EST NO. 655-NJ-1

NET CONTENTS _____

LOT NO. _____

Manufactured by

PARSONS
Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105
Office: 363 7th Avenue
New York, N.Y. 10001

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is to be used only in the formulation of anticoagulant rodenticides. Formulators using this product are responsible for providing data to support their registrations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Destroy when empty. Bury in a safe place or burn. Keep away from fumes.

180

18