

PM 04
Prentox Larva-Lur with Dylox
EPA Registration No. 655-791

655-791 11-19-97
Accepted Label on pg. 2 of 2

1 of 2

PRENTOX® LARVA-LUR with DYLOX®

An Insecticide for Control of American, German, and Oriental Cockroaches, Crickets, Ants, Pillbugs, Earwigs, and Flies

Not For Residential Indoor Use
Not For Lawn/Turf Use

ACTIVE INGREDIENT:	Dimethyl (2,2,2-Trichloro-1-Hydroxyethyl) Phosphonate	5.0%
INERT INGREDIENTS:		95.0%
	TOTAL:	100.0%

Dylox® is a Registered Trademark used under license.

EPA REG. NO. 655-791

EPA EST. NO. 655-GA-1

NET CONTENTS

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Manufactured by:

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through the skin. Do not contaminate feed or food. Avoid prolonged contact with skin. Wash thoroughly after using and before eating or smoking. May cause eye irritation. Avoid eye contact.

SYMPTOMS OF POISONING: A sense of "tightness" in chest, sweating, contracted pupils, stomach pains, vomiting, diarrhea.

Personal Protective Equipment: Applicators and other handlers must wear: Long sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Aerial application is prohibited.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

This product may cause skin sensitization in some people.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife and is extremely toxic to aquatic invertebrates. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters by cleaning of equipment or disposal of wastes.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or poison control center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: ANTIDOTE — Atropine Sulphate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine, only if symptoms of cholinesterase inhibition are present.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: LARVA-LUR is a ready-to-use, quick-acting product for the control of cockroaches, flies, crickets, Texas Harvester Ants, pillbugs, and earwigs.

User Safety Recommendations: Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2012

INDOOR USE

USE RESTRICTIONS: For control of American, German, and Oriental cockroaches and flies. For use inside of food processing plants (non-food/non-feed areas), stores, restaurants (non-food areas), and similar commercial buildings.

NOTE: Do not use in the food/feed areas of food/feed handling establishments. Do not use in edible product areas of food or feed processing plants, restaurants or other areas where food or feed is commercially prepared or processed. Do not use in serving areas while food is exposed. Applications are permitted in non-food areas of food handling establishments. Examples of non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage areas (after canning and bottling).

PRE-APPLICATION DIRECTIONS: Before applying, cover all food, feed, drinking water, and food utensils to insure that dying insects will not contaminate these items. Use a container marked to hold 4 ounces of bait but not an actual household measuring cup.

APPLICATION DIRECTIONS: Apply tablespoonful amounts of bait in wall voids, cracks and crevices, or on removable substrates such as cardboard, paper, or plastic bait stations if placed in areas inaccessible to children and pets. At conclusion of control program, remove substrates and properly dispose of uneaten baits.

IN & AROUND BUILDINGS & NON-FEED/NON-FOOD USE

USE RESTRICTIONS: For control of flies, crickets, Texas Harvester Ants, pillbugs, and earwigs. For use in and around dairy barns, milk room patios, battery poultry establishments, stables, stock pens, kennels, animal hospitals, garbage areas, and outdoor latrines. For use on patios and in picnic areas. For use in non-crop and non-feed areas.

Do not treat areas such as drinking cup, mangers, or troughs where livestock feed. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not apply bait to areas accessible to animals.

Do not contaminate water, feedstuffs, feed handling equipment, or milk handling equipment. Do not apply bait to areas accessible to animals.

PRE-APPLICATION DIRECTIONS: Before applying, cover all feed, drinking water, and feed utensils to insure that dying insects will not contaminate these items.

APPLICATION DIRECTIONS:

- 1. Hand Broadcast (in and around buildings).** Use a container marked to hold 4 ounces of bait but not an actual household measuring cup. Scatter lightly in strips, at the rate of 4 oz. per 1000 sq. ft. on floor, windowsills, and other clean horizontal areas and directly on manure and garbage, where flies breed and congregate. For faster kill, apply to a moistened surface in battery poultry establishments, bait may be applied under capes, on alley-ways, and walk-ways, but not within reach of birds.
- 2. Spreader Broadcast (around buildings).** Use spreader to broadcast at the rate of ½ to 1 pound per 1000 sq. ft. around the perimeter of homes and other buildings.
- 3. Mound Treatment (Texas Harvester Ant).** Scatter ½ to 1 cup over each mound surface. Inspect mound regularly and repeat if ant activity is spotted. Apply only by hand broadcast.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool dry place in original container, out of reach of children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag or plastic container into application equipment. Then dispose of empty bag or empty plastic container in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE

Seller's guarantee shall be limited to the terms of this label, and subject thereto, the buyer assumes any risks to persons or property rising out of the use or handling and accepts the product on these conditions.

Do not use this product for any uses other than those specified on this label.

1/97

