

# New LF-10<sup>®</sup>

## HOSPITAL DISINFECTANT CONCENTRATE

Disinfectant-Deodorant

**GERMICIDAL\*** • Staphylocidal  
• Pseudomonacidal  
• Tuberculocidal

**FUNGICIDAL\*** (Against pathogenic fungi)

**VIRUCIDAL\*** (Against Herpes simplex,  
Vaccinia, and Influenza A<sub>2</sub>  
viruses)

\*On Environmental Surfaces

ACTIVE INGREDIENTS: Potassium o-benzyl-p-chlorophenate, 9.4%; Isopropyl Alcohol, 4.2%; Sodium Dodecylbenzene sulfonate, 3.9%; Potassium o-phenylphenate, 2.9%; Tetrasodium ethylenediamine tetraacetate, 2.4%.

INERT INGREDIENTS: 77.2%

E.P.A. Registration No. 675-37-AA

### DANGER:


### KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Do not reuse empty container. Rinse thoroughly with water before discarding or returning to reconditioner.

NATIONAL LABORATORIES  
LEHN & FINK INDUSTRIAL PRODUCTS DIVISION OF  
STERLING DRUG INC.

Montvale, N.J. 07645

Plant: Toledo, Ohio 43612



### ECONOMICAL

### VERSATILE

New LF-10 Has Many Uses

#### TO DISINFECT FLOORS AND WALLS:

Use the recommended dilution of LF-10 in routine housekeeping. Just add the recommended dosage of LF-10 to water; then use for scrubbing and mopping.

#### IN THE OPERATING ROOM:

Use the recommended dilution of LF-10 as the standard disinfectant for floors, walls, equipment, and as a preliminary soak for contaminated instruments.

#### FOR CONCURRENT OR TERMINAL DISINFECTION:

Use the recommended dilution of LF-10 on furniture, floors, walls, and utensils handled by patients with contagious diseases. Use regularly in tuberculosis wards and isolation rooms.

#### IN THE NURSERY:

Use the recommended dilution of LF-10 to wipe off bassinets and equipment and to clean floors and walls. Bassinets and other equipment that may be in direct contact with infants should be rinsed following disinfection.

#### IN FOOD SERVICE:

Use the recommended dilution of LF-10 to wipe down food carts and equipment, to disinfect and deodorize garbage cans and surrounding areas and to mop floors in kitchen and dining areas. Food contact surfaces treated with LF-10 solution should be thoroughly scrubbed with detergent and rinsed with potable water before reuse.

**HOW TO USE New LF<sup>®</sup>-10 HOSPITAL DISINFECTANT CONCENTRATE**  
For Economy In Use Always Dilute LF-10 According to Directions  
One Tablespoonful Equals about 1/2 Fluid Ounce or 15 cc.

PURPOSE	Correct Solution Strength	To prepare proper dilutions in required quantities, add full strength LF-10 to water as below:	
		LF-10	Water
FOR GENERAL DISINFECTION of equipment, floors, walls, and furniture. Apply with mop, cloth, sponge, or spray equipment.			
FOR DISINFECTION OF SURGICAL AND DENTAL INSTRUMENTS: Rinse off blood and exudates. Immerse instruments in a solution of LF-10 for 20 minutes. After immersion, rinse instruments with sterile water.	1/2% (1:200)	1 tsp. (5 cc.) 2/3 oz. (20 cc.) 1-1/3 oz. (40 cc.)	1 quart 1 gallon 2 gallons
FOR DISINFECTION OF THERMOMETERS (ORAL AND RECTAL): Immerse in solution for 20 minutes. Rinse with potable water before reuse.			

#### FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites or gelatin solution; if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

#### NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, as well as oxygen, and measures to support breathing, manually or mechanically, may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

675-37