

# Wettable Sulphur

## ACTIVE INGREDIENT:

Sulphur 90%

## INERT INGREDIENTS:

10%

EPA Reg. No. 769-191 EPA Est. 49832-AL I TOTAL 100%

Sulphur - 90% through 325 mesh screen

## KEEP OUT OF REACH OF CHILDREN CAUTION

**HAZARDS TO HUMANS & DOMESTIC ANIMALS:** Harmful if swallowed. Avoid breathing dust. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not use for medical purposes or as a feed supplement.

**FIRST AID: IF IN EYES,** immediately flush with water for 15 minutes. Call a physician. Irritation persists. **FOR SKIN CONTACT,** wash with soap and water. **IF INHALED** and overcome by vapor or dust, remove immediately from exposure and call a physician. Administer artificial respiration if breathing is irregular or has stopped. **IF SWALLOWED,** drink 1 or 2 glasses of water and induce vomiting by sticking finger in back of throat. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician.

**ENVIRONMENTAL HAZARDS:** Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

**USE PRECAUTIONS:** Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality. During periods of high temperatures, sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use sulphur with oil or within 3 weeks after of application. When using Lead Arsenate with SECURITY Wettable Sulphur, always add the same amount of hydrated lime or other salver as would be used when applying Lead Arsenate without sulphur. If the crop to be treated will be used for commercial canning or freezing purposes, consult food processor before applying sulphur.

**REENTRY INTERVAL:** Do not reenter treated fields within 24 hours.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**GENERAL INFORMATION:** When more than one dosage range is given, use the higher rate when conditions for diseases are most favorable and use the lower rate when conditions for diseases are least favorable.

**CITRUS:** To control rust mite, apply 5 to 10 pounds per 100 gallons water. During cool weather, use 10 pounds SECURITY Wettable Sulphur per 100 gallons water.

**APPLES:** To control apple scab and powdery mildew, apply 6 to 8 pounds per 100 gallons water. Make application between periods of rainfall. Repeat as necessary to cover new foliage growth during pink, petal fall and cover spray periods.

**PEACHES AND PLUMS:** To control brown rot and scab, use 6 pounds per 100 gallons water. Follow locally issued spray schedule for number and frequency of applications. Usually 6 applications are made: No. 1 - when 7/8 petals have fallen; No. 2 - when 7/8 shucks have shed; No. 3 - two weeks later; No. 4 - two weeks after No. 3; No. 5 - four weeks before harvest; No. 6 - two weeks before harvest. Do not apply within 2 weeks of an oil spray treatment.

### FLC-WERS AND ORNAMENTALS

**APPLICATION:** To control the diseases listed below and spider mites, use Wettable Sulphur as a dust, or spray using 6 pounds per 100 gallons of water. Applications should be made at 5 to 7 day intervals beginning when the disease first appears and continuing until all danger from the disease is past. During periods of humid or rainy weather, it may be necessary to make applications as often as every 2 to 3 days.

**POWDERY MILDEW:** Aster, Chrysanthemum, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euponymus, Golden Fleece, Hibiscus, Honeysuckle, Hydrangea, Lady's Mantle, Ligustrum, Lilac, Clematis, Myrtle, Phlox, Poplar, Silver Vine, Spiraea, Sunflower, Verbena, Willow, and Zinnia.

**LEAF SPOT:** Calendula, Cherry Laurel, Clematis, Columbine, English Ivy, Foxglove, Hydrangea, Petunia, Phlox, Sage, Smilax and Snapdragon.

**BLACK SPOT:** Rose.

**ROSE POWDERY MILDEW:** Rose.

**BOTRYTIS BLIGHT:** Hydrangea.

**RED SPIDERS:** Arbovitae, Carnation, Cedars, Chrysanthemum, Clematis, Hydrangea, Rose, Snapdragons and Sweetpeas.

### STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state or local procedures. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residues into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

**NOTICE:** The seller makes no implied warranty of merchantability or any other warranties which extend beyond the description on this label.

387319

# SECURITY

ACCEPTED

JAN 6 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, all amended for the pesticide registered under EPA Reg. No. 769-191

SECURITY LAWN & GARDEN PRODUCTS CO.  
Manufacturers Fort Valley, Georgia