



**DIRECTIONS**

Keep can sealed between applications to prevent loss of toxicant.

For use in BARNs, MILKROOMS, MILKSHEDS, LIVESTOCK AREAS, HOG PENS, STABLES, FEEDLOTS, DOG KENNELS, FURBEARING ANIMAL UNITS, sprinkle MACKWIN SUGAR FLY BAIT lightly on floor, stalls, aisles, stanchion area, window sills, ledges, and on the ground near entrances especially to the feed storage room. Apply chiefly where flies like to congregate such as sunny areas, near water and on manure. Do not treat freshly limed floors. Apply daily 1/4 pound per 1000 square feet until fly population is reduced. Repeat as needed. MACKWIN SUGAR FLY BAIT increases contact by killing flies on animals and resting surfaces above the floor.

**POULTRY HOUSES:** Sprinkle MACKWIN SUGAR FLY BAIT on litter and floor in and about poultry houses where flies tend to congregate, but keep out of reach of poultry. Apply 1/4 pound per 1000 square feet. Repeat as needed. REMOVE MANURE FREQUENTLY.

For use in HOMES, BACKYARDS, PICNIC AREAS, PATIOS, SERVICE STATIONS, LOADING DOCKS, OUTDOOR LATRINES, apply 1/4 pound per 500 square feet of MACKWIN SUGAR FLY BAIT around these areas. Also sprinkle in and around garbage cans.

**SUGAR FLY BAIT**

Containing Vapona® Insecticide

Kills by contact or fumes, pesky, disease carrying resistant and nonresistant flies.

Sprinkle on fly feeding, breeding and resting surfaces. Use in Barns, Milkrooms, Hog Pens, Stables and other livestock areas as listed.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

See side panel for other precautions

**ACTIVE INGREDIENTS:**

2, 2-dichlorovinyl dimethyl phosphate*	0.46%
Related Compounds*	0.04%
<b>INERT INGREDIENTS:</b>	<b>99.50%</b>
	<b>Total 100.00%</b>

\*Equivalent to .5% Vapona® Insecticide (formulated with 1%)

EPA Reg. No. 995-45  
EPA Est Reg. No. 995-MN-1

**Contents 32 Oz. (2 Lbs.)**

**CAUTION:** Keep out of the reach of children. HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs, milk or milking utensils. If swallowed, induce vomiting immediately and get medical attention.

Keep out of reach of domestic animals.

**DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.**

"This product is toxic to fish, birds, and other wildlife."

"Keep out of lakes, streams, or ponds."

"Do not apply where runoff is likely to occur."

"Do not contaminate water by cleaning of equipment, or disposal of wastes."

"Apply this product only as specified on this label."

**PHYSICIAN NOTE:** Atropine is antidotal.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**NOTE:** This product is sold by weight not by volume so container may not appear to be full.

Manufactured by The MACKWIN Company Winona, Minnesota 55987





# SUGAR FLY BAIT

## Containing Vapona® Insecticide

Kills by contact or fumes, pesky, disease carrying resistant and nonresistant flies.

Sprinkle on fly feeding, breeding and resting surfaces. Use in Barns, Milk-rooms, Hog Pens, Stables and other livestock areas as listed.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:	
2, 2-dichlorovinyl dimethyl phosphate*	0.46%
Related Compounds*	0.04%
INERT INGREDIENTS:	99.50%
	Total 100.00%

\*Equivalent to .5% Vapona® Insecticide (formulated with 1%)

EPA Reg. No. 995-45  
EPA Est Reg. No. 995-MN-1

Contents 32 Oz. (2 Lbs.)

Keep can sealed between applications to prevent loss of toxicant.

For use in BARNs, MILKROOMS, MILKSHEDS, LIVESTOCK AREAS, HOG PENS, STABLES, FEEDLOTS, DOG KENNELS, FURBEARING ANIMAL UNITS, sprinkle MACKWIN SUGAR FLY BAIT lightly on floor, stalls, aisles, stanchion area, window sills, ledges, and on the ground near entrances especially to the feed storage room. Apply chiefly where flies like to congregate such as sunny areas, near water and on manure. Do not treat freshly limed floors. Apply daily 1/4 pound per 1000 square feet until fly population is reduced. Repeat as needed. MACKWIN SUGAR FLY BAIT increases contact by killing flies on animals and resting surfaces above the floor.

**POULTRY HOUSES:** Sprinkle MACKWIN SUGAR FLY BAIT on litter and floor in and about poultry houses where flies tend to congregate, but keep out of reach of poultry. Apply 1/4 pound per 1000 square feet. Repeat as needed. REMOVE MANURE FREQUENTLY.

For use in HOMES, BACKYARDS, PICNIC AREAS, PATIOS, SERVICE STATIONS, LOADING DOCKS, OUTDOOR LATRINES, apply 1/4 pound per 500 square feet of MACKWIN SUGAR FLY BAIT around these areas. Also sprinkle in and around garbage cans.

Keep out of reach of domestic animals.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

"This product is toxic to fish, birds, and other wildlife."

"Keep out of lakes, streams, or ponds."

"Do not apply where runoff is likely to occur."

"Do not contaminate water by cleaning of equipment, or disposal of wastes."

"Apply this product only as specified on this label."

**PHYSICIAN NOTE:** Atropine is antidote.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**NOTE:** This product is sold by weight not by volume so container may not appear to be full.

Manufactured by The MACKWIN Company, Winona, Minnesota 55987