

1001-78

9/6/2013

1/17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Ms Carrie M. Tackema  
Cleary Chemicals, LLC  
11901 S. Austin Ave.  
Alsip, IL 60803

SEP 06 2013

Subject: Product Name: 3336 Plus Systemic Fungicide  
EPA Reg. No. 1001-78  
Submission date: 8/13/13  
Notification for label changes per PRN 98-10: Update the registrant address and warranty statement  
Decision Number 482304

Dear Ms Tackema:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 8/13/2013 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have questions concerning this letter, please call Banza Djapao at 703-305-7269 or via email at [djapao.banza@epa.gov](mailto:djapao.banza@epa.gov), or you may call me at 703-308-3194.

Sincerely,

A handwritten signature in cursive script that reads "Shaja B. Joyner".

Shaja B. Joyner  
Product Manager 20  
Fungicide Branch  
Registration Division (7504P)

2/17

# EXPEDITE

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other:	OPP Identifier Number
--	---	---	-----------------------

## Application for Pesticide - Section I

1. Company/Product Number <b>1001-78</b>	2. EPA Product Manager <b>Tony Kish</b>	3. Proposed Classification  <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <b>3336 Plus</b>	PM# <b>22</b>	
5. Name and Address of Applicant (Include ZIP Code) <b>Cleary Chemicals, LLC.</b> <b>11901 S. Austin Ave.</b> <b>Alsip, IL 60803</b> <i>Please send all correspondence to "contact point" listed below</i>  <input checked="" type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

## Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)


### NOTIFICATION PER PRN 98-10

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**CONTACT:** Carrie M. Tackema (919) 379-2528 or [carrie.tackema@us.nufarm.com](mailto:carrie.tackema@us.nufarm.com)  
 Nufarm Americas, Inc.  
 4020 Aerial Center Parkway, Suite 101  
 Morrisville, NC 27560

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No			<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other
*Certification must be submitted		If "Yes" Unit Packaging wgt.    No. per container	If "Yes" Package wgt.    No. per container		
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container  1 GAL, 2.5 GAL		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product					
<input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Other <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

## Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name <b>Carrie M. Tackema</b>	Title <b>Regulatory Manager</b>	Telephone No. (Include Area Code) <b>(919) 379-2528</b>
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title <b>Regulatory Manager</b>	
4. Typed Name <b>Carrie M. Tackema</b>	4. Date <b>August 13, 2013</b>	



Cleary Chemical LLC  
11901 S. Austin Avenue  
Alsip, IL 60803  
Phone: 708-377-1330 Fax: 708-377-1333

8/13/13

4020 Aerial Center Parkway, Suite 101  
Morrisville, NC 27560  
Phone: 919-379-2510 Fax: 919-467-5923  
www.us.nufarm.com

Via Courier Delivery

August 13, 2013

Mr. Tony Kish, #22  
USEPA (7504P)  
Document Processing Desk (NOTIF)  
Room S4900, One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

**Subject: Notification(s) per PRN 98-10  
Spectro 90 WDG, EPA Reg. No. 1001-72; and  
3336 Plus Systemic Fungicide, EPA Reg. No. 1001-78**

Dear Mr. Kish:

On behalf of Cleary Chemical, LLC, enclosed please find two (2) Notifications for the above-reference products. The purpose of these Notifications is to update the company address and warranty statement.

Also enclosed in support of these Notification(s) are the following:


- Application(s) for Pesticide Registration for each product;
- Two (2) copies of proposed product labeling for each product (one copy with changes highlighted, and one in pdf format on CD-rom).

Should you have any questions or require additional information, please contact me at (919) 379-2528 or by email at [carrie.tackema@us.nufarm.com](mailto:carrie.tackema@us.nufarm.com).

Sincerely,

Carrie M. Tackema  
Regulatory Manager

Enclosure(s)



# 3336 PLUS™ Systemic Fungicide

9/17

[Optional marketing claims:

- With Cleartec
- With Cleartec Activation Technology
- With Activation Technology]

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

**NOT FOR HOMEOWNER USE**

**Active Ingredient:**

Thiophanate-methyl (dimethyl-4,4'-o-phenylenebis[3-thioallophanate])* .....	19.4%
Other Ingredients .....	80.6%
<b>Total:</b>	<b>100.0%</b>

\*This product contains 2.0 lbs thiophanate methyl per gallon.

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**  
 See side panels for additional precautionary statements

**FIRST AID**

<b>If On Skin Or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor for treatment advice.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If In Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<p><b>For emergency information, call (732) 329-8399 or your poison control center.</b></p> <p><b>For medical and transportation emergencies only call 24 hours a day CHEMTREC (800) 424-9300.</b></p> <p><b>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</b></p>	
<p><i>See side/back panels for additional precautionary statements</i></p>	

EPA Reg No. 1001-78

Net Contents: 1 Quart, 1 Gallon, 2.5 Gallon

EPA Establishment No.

**Manufactured For:**  
**CLEARY CHEMICAL LLC**  
**11901 S. Austin Avenue**  
**Alsip, IL 60803**

S/17

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Do not get on skin or clothing. Avoid contact with eyes.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are made of any waterproof materials. If you want more options, follow the instructions for Category C on an EPA chemical resistance selection chart.

Handlers mixing, loading and applying the product as a dip must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant apron
- For overhead exposure wear Chemical-resistant headgear
- When cleaning equipment, wear a Chemical-resistant apron

All other mixers, loaders and applicators must wear:

- Coveralls over short sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves (such as or made out of any waterproof material, selection category C)
- Chemical-resistant apron for mixers, loaders and other handlers exposed to concentrate, and for cleaning equipment.
- For overhead exposure wear Chemical-resistant headgear

**USER SAFETY REQUIREMENTS**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

**DIRECTIONS FOR USE:**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for over head exposures.

Exemption: if the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter treated areas if there will be no contact with anything that has been treated.

**Non-Agricultural Use Requirements**

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Do not enter or allow others to enter until sprays have dried.

**GENERAL INFORMATION**

**Failure to follow directions and precautions on this label may result in crop injury, poor disease control, and/or illegal residues.**

3336 PLUS is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 PLUS with ground or overhead equipment, using sufficient volume of spray to provide thorough coverage. Do not apply with fixed wing or rotary aircraft. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. This product is not for homeowner use, and can only be used by individuals/firms licensed under the direction of or registered by the state to apply ornamental or turf pest control products. Chemigation instructions follow "Directions For Use". Read and follow these instructions carefully for this method of application.

7/17

**Resistance Management:** To avoid the development of tolerant strains of fungi, 3336 PLUS should be used with fungicides with different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 PLUS. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using PLUS as recommended, and the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 PLUS can remain useful for disease control.

**Mixing Instructions:** SHAKE WELL BEFORE USING: Some settling may occur during prolonged periods on non-use. High pH environments cause a shortened tank life for diluted product. The buffering of tank water to pH 6-7 prior to the addition of 3336 PLUS is recommended. Add required amount 3336 PLUS to partially filled tank (1/2 total volume), agitate by mechanical or hydraulic means, add tank mix product if used (see below), agitate again and then add remaining required amount of water. Continuous agitation is recommended to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

**Tank Mixing Instructions:** 3336 PLUS may be compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. 3336 PLUS may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

## CHEMIGATION REQUIREMENTS

### Generic Requirements

1. Apply this product only through the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move; flood (basin); or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### Specific Requirements

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be

discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

- 3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
- 4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Non-Specific Requirements**

- 1. Remove scale, pesticide residue, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.
- 2. Prepare a suspension of product in the mix tank or stock bucket. Fill the tank with 1/2 or 3/4 of the desired amount of water. Start agitation and add the required amount of product to the solution along with the remaining volume of water.
- 3. Maintain a gentle agitation in the mix tank during application to assure a uniform suspension. Follow mixing instructions and tank mixing instructions previously indicated.
- 4. Start system and then uniformly inject the suspension of 3336 PLUS into the irrigation line so as to deliver the desired rate per acre. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation system.
- 5. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.
- 6. The suspension of 3336 PLUS should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

**Sprinkler (Overhead) Chemigation**

Observe all instructions in the Generic, Specific and Non-Specific Requirements sections above and the following additional requirements:

- 1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Set sprinkler system to deliver 1/10 to 1/4 inches of water per acre. Volumes of water higher than this may reduce efficacy. Application of more than recommended quantities of irrigation water per acre may result in decreased product performance. Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result. When system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained in a center pivot system, block the nozzle set nearest the well pivot injection unit to prevent spray being applied to this area. Allow sufficient time for pesticides to be flushed through all lines and all nozzles before turning off irrigation water.



9/17

**Flood (Basin) Chemigation**

1. Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
2. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements: Observe all instructions in the Generic, Specific and Non-Specific Requirements sections above and items 1 and 2 of the sprinkler irrigation requirements.

**Drip (Trickle) Chemigation**

Observe all instructions in the Generic, Specific and Non-Specific Requirements sections above.

**TURF APPLICATIONS**

3336 PLUS may be used on all fine turf applications such as commercial, residential and public (such as, home lawns, parks, athletic fields, schools, and day care centers), and golf courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermuda grass, Fescue, Ryegrass, St. Augustine grass, Zoysia grass, or their mixtures. 3336 PLUS is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 PLUS is to be used for the prevention and control of the diseases mentioned below. It has both preventative and curative activity. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod.

**Application Instructions:** Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation systems. Spray uniformly over the area to be treated. Apply recommended amounts in sufficient water to obtain thorough coverage of treatment area (2-4 gal per 1,000 sq ft is suggested). Use the highest recommended rate under conditions of severe disease pressure. For best results, apply after mowing or avoid mowing twelve hours after application. For root pathogens, lightly water the treatment area to move the fungicide into the active root zone with one to two tenths inch of water. Excessive irrigation may move application below active root zone and reduce application effectiveness. Green design and drainage will influence irrigation practices. When tank mixing with contact action fungicides for foliar diseases, applications should be allowed to dry on leaf surfaces. Normal watering may proceed after sprays have dried.

**Table 1: Maximum Individual Application Rates and Minimum Re-Treatment Intervals**

Do not exceed the amounts per acre or reduce the re-treatment interval indicated below:

Use Site	Maximum application rate of 3336 PLUS	Minimum Re-Treatment Interval	Comments
Residential or Public Areas	1.36 Gallons/Acre (4 oz / 1,000 sq.ft.)	14 days	
Golf Course Tees, Greens, Aprons	4.08 Gallons/Acre (8 oz / 1,000 sq.ft.)	14 days	
Golf Course Fairways –except Florida	2.73 Gallons/Acre (8 oz / 1,000 sq.ft.)	14 days	Excludes Florida
Golf Course Fairways – Florida Only	1.36 Gallons/Acre (4 oz / 1,000 sq.ft.)	14 days	Florida Only One Application Only During Overseeding

10/17

**Table 2: Maximum Seasonal Application Rates**

Do not exceed the following amounts of product per Acre per season:

Use Site	Maximum Gallons 3336 PLUS per Acre per Season	Fluid Ounces 3336 PLUS per 1,000 sq.ft.	Comments
Residential or Public Areas	5.44 Gallons	16 fl.oz.	4 Apps @ 2-4 oz
Golf Course Tees, Greens, Aprons	10.88 Gallons	32 fl.oz.	
Golf Course Fairways –except Florida	2.73 Gallons	8 fl.oz.	1 App @ 8 oz, or 2 Apps @ 4 oz, or 4 Apps @ 2 oz
Golf Course Fairways – Florida Only	1.36 Gallons	4 fl.oz.	Florida Only – During Overseeding 1 App @ 4 oz, or 2 Apps @ 2 oz

Disease(s) Controlled	Rate of 3336 PLUS fl oz/1,000 sq ft	Remarks
<b>Anthracnose, basal</b> <i>Colletotrichum graminicola</i> <b>Anthracnose, foliar</b> <i>Colletotrichum graminicola</i>	4-8 2-4	For prevention in historic areas of disease pressure, apply twice at 14 day (d) intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears and continue at 14-28 d intervals. Rotations and/or tank mix combinations with chlorothalonil or triadimefon can be utilized.
<b>Bermudagrass Decline</b> <i>Gaeumannomyces graminis</i> var. <i>graminis</i> <b>Take-All-Patch</b> <i>Gaeumannomyces graminis</i> var. <i>avenae</i>	4-8	Apply in mid-July or when disease symptoms first appear and repeat at 14-28 d intervals for suppression. Use higher rates under most severe disease expression. Water treatment into active root zone. Follow proper agronomic recommendations to maintain plant vigor.
<b>Cool Season Brown Patch</b> <i>Rhizoctonia cerealis</i> <b>Necrotic Ring Spot</b> <i>Leptosphaeria korrea</i>	4-8	For prevention, apply in Fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early Spring and continue at 14-28 d intervals. Water treatment into active root zone.

11/17

<b>Coprinus Snow Mold</b> <i>Coprinus psychromorbidus</i>	4-8	Apply 2 treatments at 21 d intervals in late Fall to early Winter, with the last application made just prior to first permanent snow cover. Rotations and/or tank mix combinations with PCNB can be utilized.
<b>Dollar Spot</b> <i>Moellerodiscus, Lanzia</i> <b>Large Brown Patch</b> <i>Rhizoctonia solani</i> <b>Ascochyta Leaf Blight</b> <i>Ascochyta</i> <b>Copper Spot</b> <i>Gloeocercospora sorghi</i> <b>Fusarium Patch</b> <i>Fusarium nivale</i> <b>Red Thread</b> <i>Corticium fuciforme</i> <b>Zoysia Patch</b> <i>Rhizoctonia solani</i>	2-4	Apply when disease first appears and continue at 14-28 d intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect DF) can be utilized.
<b>Fusarium Blight</b> <i>Fusarium roseum, F. triticum</i>	4-8	Apply when disease first appears at 14-28 d intervals.
<b>Gray Leaf Spot (Blast)</b> <i>Pyricularia grisea</i>	4-8	Apply preventative application before expected period of disease development. Continue applications at 14-28 d intervals.
<b>Leaf Spot</b> <i>Drechslera</i> <b>Leaf, Crown, and Root Diseases</b> <i>Bipolaris, Curvularia, Exserohilum</i>	4-8	Apply when disease first appears and make applications at 14-28 d intervals as needed. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect DF) are recommended under severe conditions.
<b>Pink Snow Mold</b> <i>Microdochium nivale</i>	2-4	Apply in late Fall to early Winter before turf has stopped all growth activity. A second application may be used in combination with chlorothalonil, PCNB, or thiram (Spotrete) at recommended rates before snow cover or during Spring thaw.
<b>Rusts</b> <i>Puccinia, Uromyces</i>	4-8	Apply at 14-28 d intervals when disease first appears. Rotations and/or tank mix combinations with chlorothalonil or mancozeb (Protect DF) are recommended.
<b>Stripe Smut</b> <i>Ustilago striiformis</i>	4-8	Apply at 14-28 d intervals when disease first appears. For prevention, apply in spring and fall.
<b>Summer Patch</b> <i>Magnaporthe poae</i>	4-8	For prevention, apply 3 applications starting late April or early May using 21 d intervals. Rotations and/or tank mix combinations may be used as part of the three application program. For suppression, apply at 14-28 d intervals when disease first appears. Water treatment into active root zone.

<b>Bentgrass Dead Spot</b>	4-6	For prevention, apply in early June or based upon local extension service recommendations. Apply at 14 day intervals. Rotations and/or tank mix combinations may be used for season long disease prevention
----------------------------	-----	---

## HORTICULTURAL APPLICATIONS

**Nursery, Greenhouse, Landscape & Interiorscape**  
**Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs**

Do not use fruits, nuts or sap from treated trees as food or feed.

3336 PLUS is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 PLUS is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 14-28 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. 3336 PLUS may be applied as a ground application using hand held, mechanical or motorized spray equipment, or as a chemigation spray, or through an applicable sprinkler irrigation system. See specific instructions below.

**Note:** The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (*Plectranthus australis*), Boston Fern (*Nephrolepis exhalta*), and Easter Cactus (*Hatiora gaertneri*).

**Application Instructions:** Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation, flood, or drip systems. Begin applications when disease first appears and repeat at 14-28 day intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix 8-24 fl oz of 3336 PLUS per 100 gal water (0.5-1.5 teaspoons per gal) and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Spray volume may range up to 400 gallons of finished spray per acre depending upon plant species and plant growth stage. For applications through irrigation systems, refer to aerial use rates indicated in the foliar application chart. For small volume applications less than 100 gallons, divide recommended rate by 16 to get the number of teaspoons of 3336 PLUS per gal.

### **Special Instructions for Proportional Injectors (e.g. Dosatron, Dosmatic, Anderson and similar equipment)**

Determine the treatment rate for each crop and pathogen from the foliar application table below. Determine the injection ratio for the individual system to be used for application. For systems using a 1:100 ratio, measure and add the exact amount of recommended material per 100 gallons to each gallon of water in a stock bucket or tank. For systems using a 1:200 ratio, multiply the recommended amount per 100 gallons by 2. For systems using a 1:50 ratio, divide the recommended amount per 100 gallons by 2. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time. An injection ratio of 1:100 is recommended for most greenhouse and nursery systems.

13/17

**FOLIAR APPLICATION: \***

Diseases Controlled	Rate of 3336 PLUS	Remarks
<b>Anthracnose</b> <i>Colletotrichum</i>	12-16 fl oz/100 gallon	Apply as buds break or at first sign of disease. Repeat at 14-28 day (d) intervals as needed during disease period.
<b>Black Spot of Rose</b> <i>Diplocarpon rosae</i>	12-16 fl oz/100 gallon	Apply early summer or at first sign of disease. Repeat every 14-28 d as needed during disease period.
<b>Brown Rot and Blight</b> <i>Monilinia, Sclerotinia, Whetzellinia</i>	12-16 fl oz/100 gallon	Apply late Spring or at first sign of disease. Repeat every 14-28 d as needed during disease period.
<b>Fusicladium and Venturia Leaf Scabs on:</b> Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	12-16 fl oz/100 gallon	Apply as buds break. Repeat every 14-28 d during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with mancozeb (Protect DF), chlorothalonil or propiconazole can be utilized.
<b>Leaf Spots and Blights caused by:</b> <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Cocomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Physalospora, Schizothyrium, Septoria, Sphaceloma</i>	12-16 fl oz/100 gallon	Apply when disease symptoms first appear. Repeat every 14-28 d during disease period. Rotations and/or tank mix combinations with mancozeb (Protect DF) or chlorothalonil can be utilized.
<b>Ovulinia Blight</b>	8-16 fl oz/100 gallon	Apply as flowers open. Repeat every 14-28 d during disease period.
<b>Powdery Mildews</b> <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	12-24 fl oz/100 gallon	Apply when disease first appears and repeat every 14-28 d during disease period. Rotations and/or tank mix combinations with mancozeb (Protect DF) or triadimefon can be utilized.
<b>Rust Diseases caused by:</b> <i>Puccinia, Gymnosporangium, Uromyces</i>	12-16 fl oz/100 gallon	Apply late Spring or when symptoms first appear. Repeat every 14-28 d during disease period. Rotations and/or tank mix combinations with mancozeb (Protect DF) or chlorothalonil are recommended.
<b>Tip Blight of Pine</b> <i>Sphaeropsis sapinea, Diplodia pinea</i>	16-24 fl oz/100 gallon	Begin application in spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 10-14 d later. Thorough coverage is essential for optimal disease control.
<b>Twig Blights, Cankers, and Diebacks</b> <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	16-24 fl oz/100 gallon	Apply when symptoms first appear. Repeat every 14-28 d during disease period.

\* Do not exceed 0.5 lbs ai per acre per application for cut flowers equivalent to 8 fl oz per 100 gallon spray, applied at the rate of 400 gallons spray per acre. Do not exceed 3.0 lbs ai per acre per application for cut flowers equivalent to 48 fl oz per 100 gallon spray, applied at the rate of 400 gallons spray per acre.

**SOIL DRENCH APPLICATION:**

Diseases Controlled	Rate of -3336 PLUS	Remarks
<p><b>Stem, Crown, and Root Rots</b> caused by:  <i>Botrytis, Cythindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia</i></p> <p><b>Black Root Rot</b>  <i>Thielaviopsis</i></p>	<p>8-16                      fl oz/100 gal</p>	<p>Apply as a drench or directed spray using hand held, mechanical or motorized spray equipment or as a chemigation drench or directed spray using applicable sprinkler irrigation systems after seeding or sticking of cuttings (8 fl oz) or after transplanting (12-16 fl oz) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is 0.25 –3.0 pints of finished mixture per sq ft depending on the media type and depth (about 4 fl oz per 4 inch pot or 8 fl oz per 6 inch pot). Repeat every 21-28 d for adequate crop protection. Note: 3336 PLUS does not control <i>Pythium</i> or <i>Phytophthora</i>. Tank mix combinations with metalaxyl, mefenoxam, etridiazole, propamocarb, or fosetyl-AI are required for the control of <i>Pythium</i> and <i>Phytophthora</i>. Do not exceed a maximum seasonal rate of 300 lb ai per acre.</p>

**PLANT DIP APPLICATION:\***

Diseases Controlled	Rate of -3336 PLUS	Remarks
<p><b>Plant or Cutting Diseases</b> caused by:  <i>Botrytis, Cythindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i></p>	<p>16-24                      fl oz/100 gal</p>	<p>Immerse plants or cuttings for 10-15 min. Remove and allow to drain.  <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.</p>
<p><b>Bulb, Corm, and Rhizome Rots</b> caused by:  <i>Botrytis, Cythindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i></p>	<p>16-24                      fl oz/100 gal</p>	<p>Soak cleaned bulbs for 15-30 min in warm solution (80-85° F). For storage disease prevention, treat bulbs preferably within 48 h after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured.  <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.</p>

\* Do not exceed 0.7 lbs ai per acre per application for dip treatment (Bulb, Cutting, Preplant, Post-thinning) or Dip tank.

15/17

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in a cool dry place.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to the label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

**Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S

16/17


TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

RV090711

3336 PLUS is a trademark of Cleary Chemicals LLC.

All other trademarks are the property of their respective owners.





**LABEL HISTORY**

<b>File Name</b>	<b>Revision Date</b>
001001-00078.draft080813	RV090710

cccccc      ccc  
c              cc  
c              c  
c              c  
c              c  
cccccc      ccc  
c              cc  
c              c  
c              c  
cccc        cccc  
c              cc  
cccccc      c  
cccccc      c  
cccccc      c