GARDEN SPRAY

Contains - Malathion, Methoxychlor and Kelthane*

CAUTION: KEEP OUT OF REACH OF CHILDREN - SEE REAR PANEL FOR ADDITIONAL CAUTIONS

USDA Reg. No. 1159-114

ACTIVE INGREDIENTS:

*Malathion	.15.00%
1. 1-bis (chlorophenyl) 2, 2, 2-Trichloroethano	1 5.00%
**Methoxychlor Technical	. 7.50%
Aromatic Petroleum Solvent	
INERT INGREDIENTS:	. 8.50%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

**7.5% Methoxychlor technical is equivalent to 6.6% 2, 2-bis (p-methoxyphenyl) 1, 1, 1-trichloroethane and 0.9% other isomers and reaction products.

NET CONTENTS ONE PINT

TWIN LIGHT GARDEN SPRAY FOR INSECT CONTROL ON SHRUBS, EVERGREENS, TREES, FRUITS AND VEGETABLES

1

DIRECTIONS plants thoroughly, repeat as necessary. iers, Kose Siugs, Fied Beetles, Inribs, Iarnished Plant Bugs, Four Lined Leat Bugs, Whiterly, Asiatic Garden Beetles and Black Blister Beetles.—3 teaspoonsful on ROSES, GLADIOLUS, SWEEL PEAS, DAHLIAS, ASTERS and CHRYSANTHEMUMS. SHRUBS and EVERGREENS: Japanese Beetles, Aphids, Spider Mites, Lacebugs.—3 teaspoonsful. Azalea Scale, Scurfy Scale, Eucoymus Scale, Soft Brown Scale, Bagworm and Boxwood Leaf Miners.—4 tea-spoonsful on AZALEAS, RHODODENDRONS, ARBORVITAE, ULLIPERS, HYDRANGEAS, EUONYMUS, PIERIS JAPONICA, LILAC and BOXWOOD. NOTE: Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula. Do not apply on Petunias on Petunias.

FRUITS: Aphids, Codling Moth, Curculio, European Red Mites, Clover Mites, Two Spotted Mites, Williamette Mites, Pear Psylla, Japanese Beetles, Black Cherry Aphids and Tent Caterpillars---4 tea-spoonsful on <u>APPLES, PEARS, PEACHES</u>. For control of the mites as listed above on <u>CHERRIES</u> and <u>PLUMS</u>, apply when mites appear in numbers. If repeat application is necessary wait 30 days to avoid excessive residue. FRUIT SPRAY SCHEDULE for APPLES, PEARS and PEACHES. Apply at petal fall and every 10 to 14 days thereafter as long as local conditions warrant. NOTE: Do not apply within 7 days of harvest on Apples, Pears, Cherries and Plums-21 days of harvest on Peaches. Injury may occur to McIntosh and Cortland varieties of Apples if applications made within 4 weeks of harvest.

Peach Tree Borer on Peach Trees-2 tablespoonsful. Spray trunks and lower branches of trees. Apply at 2 week intervals during periods of Moth flight in late Summer. Avoid spraying foliage or fruit. TREES: Spider Mites, Birch Leaf Miners, Holly Leaf Miners, European Pine Shoot Moth, Tent Cater-piliars and Aphids-4 teaspoonsful on PINE, BIRCHES, HOLLY, OAKS, MAGNOLIAS, MAPLES and SYCAMORES.

VEGETABLES: Eruitworms, Hornworms, Anhids, Spider Mites, Flea Beetles, Colorado Potato Beetles. 4 teaspoonsful on TOMATOES and PEPPERS. Cucumber Beetles. Squash Vine Borer on CUCUMBERS, MELONS 201 SQUASH. Mexican Bean Beetles, Leaf Hoppers and Corn Earworms on BEANS. NUTE: D. not apply within 7 days of harvest on the vegetables listed. Do not feed treated bean vines to meat or dairy animals. CLOVER MITES: 1 Tablespoonful. Spray lower foundations of house as well as ground, lawns and ornamental plants in an area 10 feet wide along the side of the house. Repeat as necessary. DO NOT USE INSIDE DWELLING.

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. In case of skin contact, wash with soap and water. Avoid storage near feed and foodstuffs. KEEP OUT OF **REACH OF CHILDREN**

FISH AND WILDLIFE CAUTION: This product is toxic to fish. Keep out of lakes, streams or ponds. SEACOAST LABORATORIES, INC.

257 HIGHWAY 18 • EAST BRUNSWICK, N. J. 08816

ACCE

DILUTION NOTE: 1 teaspoonful per gallon equivalent to 1 pint per 100 gallons of water. All dosages shown represent the quantity of WONDERLAWN LIQUID MULTI PURPOSE per gallon of water. Spray

The rate of dilutions apply to sprayers attached to garden hose as well as to conventional sprayers. ROSES, FLOWERS: Aphids, Spider_Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rose lers, Rose Slugs, Flea Beetles, Thrips, Tarnished Plant Bugs, Four Linzd Leaf Bugs, Whitefly, Asiatic