



TWIN LIGHT Methoxychlor 5% Granular in the Midwest and Northeast is recommended for use as a mosquito larvicide, applied as a pre-hatch winter treatment for control of "spring brood" mosquito larvae and as a surface treatment to water breeding sites when larvae are present during the spring and summer.

## DIRECTIONS

Apply at the following rates by aircraft or with a crank-operated spreader similar to that used for spreading seed and fertilizer. Apply only to known mosquito breeding sites.

Pre-hatch-make a single application during winter and early spring at the rate of 10 to 20 lbs. per acre (0.5 to 1 lb. actual methoxychlor).

Larvicide-make one or more applications to the source as needed for succeeding broods during the spring and summer. Apply at the rate of 3.5 to 10 lbs. per acre (0.13 to 0.5 lb. actual methoxychlor).

Jul 2, 1968

1159-143

NOTE: This product is toxic to fish. Keep out of lakes, streams or ponds. Shrimp and crab will be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply to water drainage areas where runoff or flooding will contaminate ponds, lakes, or streams. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Do not re-use container. Crush and bury when empty. Wash thoroughly after handling.