

100%

# PURE-GRO BOTRAN® 6 DUST

**ACTIVE INGREDIENTS:**

2,6-Dichloro-4-nitroaniline*	6%
INERT INGREDIENTS	94%
	<hr/> 100%

\*Tuco Products Company — Division of the Upjohn Company  
\*U.S. Pat. No. 3,119,736

E.P.A. Reg. No. 1202-273-AA  
E.P.A. Est. No. 11369-CA-1

**ACCEPTED**  
**SEP 25 1984**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered for the pesticide control under Reg. No. 1202-273

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

In case of eye contact, flush eyes immediately with plenty of water. Get medical attention.

**HAZARD TO HUMANS & DOMESTIC ANIMALS**

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Do not apply this product in such a manner as to directly or through drift

expose workers or other persons. The area being treated must be vacated by unprotected persons.

**ENVIRONMENTAL HAZARD**

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, ponds and streams. Do not apply when weather conditions favor drift or runoff from target area.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER" "Area to be treated with Botran on (date of

application). Do not enter without appropriate protective clothing until dusts have settled. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid and medical attention recommendations."

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of the smoke.

**NECTARINE:** For the control of brown rot blossom blight (*Monilinia*) apply 60 lbs. per acre at pink bud and full bloom. For the control of brown rot fruit decay (*Monilinia*) apply at 60 lbs. per acre. Apply 18, 10 and 1 day before harvest. Do not apply later than one day before harvest. Do not allow animals to graze treated areas.

**PEACHES:** For control of brown rot blossom blight (*Monilinia*) apply 60 lbs. per acre. Apply at 18, 10 and 1 day before harvest. For the control of rhizopus fruit rot apply 60 lbs. per acre. Apply one day before harvest. Do not apply later than one day before harvest. Do not allow animals to graze treated areas.

**SWEET CHERRY:** For the control of brown rot blossom blight (*Monilinia*) apply 50 lbs. per acre. Apply at popcorn, bloom, full bloom and petal fall. For the control of fruit decay (*Monilinia*) and rhizopus fruit rot apply 50 lbs. per acre. Apply 10 days and 1 day prior to harvest. Do not apply later than one day before harvest. Do not allow animals to graze treated area.

**GRAPES:** For the control of botrytis rot apply 30 lbs. per acre. Begin application while clusters are still loose and repeat application at two week intervals. Do not apply later than one day before harvest.

**TOMATOES:** (Greenhouse). For the control of stem canker (*botrytis*) apply 30 lbs. per acre. Dust stem of plant from ground level up to a height of 18 to 24 inches. Repeat weekly or as necessary.

**ONIONS, GARLIC: (Not for use in North Central States):** For control of white rot (*sclerotium*) apply 250-400 lbs. per acre. For direct seeding use mix into top 1.5 inches of soil one to two weeks before onions are seeded, for planting sets or buds apply broadcast and mix into top 1.5 inches, or apply same rate/43 560 linear feet of furrow to soil around sets or buds. Do not plant spinach as a follow-up crop on treated soil. When lettuce is to be planted as a follow-up crop, plow soil to 8 inches and cross disk the area prior to seeding.

**LETTUCE:** (Leaf, Greenhouse Only). For control of blight (*botrytis* rot) apply 30 lbs. per acre. Apply 7 days after transplanting. Repeating when plants are half mature. Do not apply to wilted plants or seedlings. Do not apply later than 14 days before harvest.

**LETTUCE:** (Head, Field). For the control of botrytis rot and sclerotinia wilt apply 133 pounds per acre. Apply at thinning, repeat seven to ten days later. Do not apply within 14 days of harvest.

**ROSE, HYDRANGEA:** For the control of botrytis gray mold, apply 30 lbs. per acre. Field—Apply BOTRAN 6 DUST prior to lifting for shipment or storage.

**CHRYSANTHEMUM, GERANIUM:** For the control of botrytis gray mold, apply 30 lbs. per acre. Greenhouse plants—Begin when disease is anticipated or first appears. Dust foliage and flowers at 7 to 14 day intervals. **Stock Cuttings:** Apply 8½ to 16½ lbs. as a uniform dust to cuttings and shipping cartons or cuttings in propagating bench, shipment or storage.

**GLADIOLUS:** For control of stromatium dry rot apply 250-265 lbs. per 14,500 linear feet or row. Apply in a 12-inch band centered on furrow at time of planting. For control of botrytis blight apply 30 lbs. to bunched cut flower spikes. From one acre after harvesting and before storage. Use sufficient pressure to insure coverage between spikes.

### PUREGRO CONDITIONS OF SALE AND WARRANTY

PUREGRO COMPANY warrants only that the product conforms to the chemical description on the label and is reasonably fit for the use stated on the label when used in strict accordance with the directions under normal conditions of use.

Critical and unforeseen factors beyond PUREGRO COMPANY'S control prevent it from eliminating all risks in connection with the use of the product. Such risks include, but are not limited to, damage to plants and crops to which the product is applied, lack of complete product control by PUREGRO COMPANY and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein and even though label directions are followed.

Therefore, this warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseen to PUREGRO COMPANY. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this product.

PUREGRO COMPANY makes no other warranty, express or implied, including any other express or implied warranty of fitness for a particular purpose or of merchantability. In no case shall PUREGRO COMPANY be liable for consequential, special, or indirect damages resulting from the use or handling of this product. PUREGRO COMPANY does not authorize any agent or salesman to make any other warranty, guarantee or representation, express or implied, concerning this product, except those expressly set forth on this product label, and denies any such warranty, guarantee or representation, express or implied, made by any agent or employee of PUREGRO COMPANY, except those expressly set forth in this product label. The foregoing conditions of sale and warranty can be varied only by an agreement in writing signed by a fully authorized representative of PUREGRO COMPANY.

